

Left pleural effusion in a young woman with genital tuberculosis

Cristina Albrici^{1,2}, Jacopo Cefalo^{1,2}, Michele Mondoni^{1,2}, Alessia Moro³, Manuela Bimbatti³, Umberto Gianelli^{2,3}, Stefano Centanni^{1,2}

¹Respiratory Unit, ASST Santi Paolo e Carlo, Milan; ²Department of Health Sciences, University of Milan; ³Pathology Unit, ASST Santi Paolo e Carlo, Milan, Italy

Abstract

Disseminated tuberculosis is a rare form of tuberculosis that can cause severe illness if diagnosed and treated late. We present the case of a young Senegalese woman who had a miscarriage due

to a pelvic inflammatory disease, followed by the development of a left pleural effusion. Despite laparoscopic findings and a salpinx biopsy that revealed necrotizing granulomas, only microbiological examinations of pleural biopsies revealed the final diagnosis of disseminated, drug-sensitive tuberculosis.

Introduction

Tuberculosis (TB) is one of the leading infectious diseases and causes of death worldwide [1]. The optimal management of disease relies on a fast microbiological diagnosis and drug resistance detection with prompt initiation of effective pharmacological treatment [2].

Disseminated TB, which is a rare form of the disease due to hematogenous or lymphatic spread of *Mycobacterium tuberculosis complex* (MTB) from the primary site of infection, may be characterized, in high-income countries, by atypical or rare clinical pictures of extrapulmonary disease (e.g., genital TB) [3-5]. Poor knowledge of this form of the disease may result in a delayed diagnosis and therapy with subsequent spread of MTB to multiple organs and potentially severe disease [3-5].

Case Report

A 27-year-old Senegalese woman, living in Italy for two years and with a negative past medical history, was referred to our Pulmonology Department for a left pleural effusion.

One month before, following a miscarriage complicated by bilateral salpingitis, she underwent a laparoscopic left salpingectomy. During surgery, hyperemic and edematous salpingus, free abdominal effusion and several whitish, miliary-shaped peritoneal lesions, involving diaphragm and visceral serous, were detected. Abdominal fluid was sent for microbiological investigations, peritoneal and salpinx biopsies were collected only for histopathological analysis. All the microbiological investigations, including smear microscopy, culture and nucleic acid amplification tests (NAAT) for MTB were negative. Following the diagnosis of pelvic inflammatory disease (PID), oral antibiotic therapy with doxycycline and metronidazole were prescribed and the patient was discharged home without any symptom. Histopathological analysis showed chronic necrotizing granulomatous inflammation with multinucleated giant cells (Figure 1A); histochemical evaluation by Ziehl-Neelsen stain resulted negative.

One month after the discharge, due to the onset of exertional dyspnea, without cough, fever and malaise, the patient underwent a chest X-ray and then computed tomography (CT) scan and was admitted to our Pulmonology Unit.

The CT scan showed a large left pleural effusion inducing

Correspondence: Cristina Albrici, Respiratory Unit, ASST Santi Paolo e Carlo, Department of Health Sciences, University of Milan, Via Di Rudini 8, 20142 Milano, Italy.
Tel. +39.02.81843025 - Fax: +39.02.81843037.
E-mail: cristina.albrici@unimi.it

Key words: disseminated tuberculosis; genital TB; medical thoracoscopy; XpertMTB/RIF; pleural biopsy.

Contributions: AC, JC, conception and design, manuscript drafting; MM, contribution to manuscript drafting; ACM, UG, SC, critically revising of the manuscript for important intellectual content. All the authors read and approved the final version of the manuscript and agreed to be accountable for all aspects of the work.

Conflict of interest: none of the authors has a conflict of interest to disclose.

Ethics approval and consent to participate: no Ethical Committee approval was required for this case report by the Department because this article does not contain any studies with human participants or animals. Informed consent was obtained from the patient included in this study.

Consent for publication: the patient gave her written consent to use her personal data for the publication of this case report and any accompanying images.

Received: 3 October 2022.
Accepted: 3 November 2022.
Early view: 14 November 2022.

Publisher's note: all claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

©Copyright: the Author(s), 2022
Licensee PAGEPress, Italy
Monaldi Archives for Chest Disease 2023; 93:2450
doi: 10.4081/monaldi.2022.2450

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial International License (CC BY-NC 4.0) which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

dysventilation of the underlying lung parenchyma, associated with bilateral pleural thickenings and a minimal pericardial effusion. No lung parenchymal abnormalities were detected (Figure 2). Chest ultrasound confirmed the presence of left free pleural effusion. No homolateral mammary lymph node was detected. No previous thoracic radiological investigations were performed by the patient.

Blood tests showed increased C-reactive protein values (2.42 milligrams (mg)/dL; normal values: 0.03-0.5 mg/dL) without leukocytosis (white blood cells: 4450/uL, normal reference range: 3600-9200/uL). Human immunodeficiency virus (HIV) antibodies 1 and 2 were negative.

In the suspicion of tuberculosis, in the absence of sputum, and in the presence of a large pleural effusion that could have hidden abnormalities of the underlying lung parenchyma, a bronchoalveolar lavage (BAL) was performed; smear microscopy, XpertMTB/RIF and culture resulted negative. A medical thoracoscopy was then performed. The endoscopic pattern was characterized by markedly hyperemic visceral, parietal, diaphragmatic and pericardial pleura with diffuse and confluent whitish nodules (Figure 2). Microbiological investigations on aspirated pleural fluid (800 mL) were all negative; smear microscopy on pleural biopsies resulted positive and XpertMTB/RIF was positive for rifampicin-susceptible MTB, subsequently confirmed by liquid culture results. Drug susceptibility testing was performed and no drug resistance to the first-line anti-TB drugs was detected. Histopathological examinations showed necrotizing granulomas (Figure 1 B-D). NAAT was then performed on salpinx biopsies

and resulted positive. A diagnosis of disseminated tuberculosis was made.

The standardized regimen recommended by the World Health Organization (*i.e.*, isoniazid 300mg/day, rifampicin 600mg/day, ethambutol 1600mg/day, and pyrazinamide 1500mg/day) was administered for a 2-month intensive therapy, which was followed by a 4-month continuation phase with isoniazid and rifampicin. Her clinical recovery was prompt and chest X-ray performed two months after did not detect any pulmonary abnormality and pleural effusion (Figure 3C).

Discussion

Tuberculosis is an important clinical and public health issue worldwide, causing 1.2 million TB deaths among HIV-negative people and an additional 208,000 deaths among HIV-positive people in 2020. It is estimated 10 million people were newly diagnosed with TB in 2019 [1].

According to the World Health Organization (WHO), Italy has a low incidence of the disease (<20/100,000), with a clear prevalence of importation cases [6].

Disseminated TB is a rare form of the disease (1-5% of all TB cases) defined by MTB identified in blood or bone marrow samples, from liver biopsies, or from specimens collected from two or more non-contiguous organs [3-5]. Its pathogenesis, largely unknown, is determined by the hematogenous or lymphatic spread




Figure 1. A) Salpinx biopsy showing giant cells and granulomas (Hematoxylin & Eosin stain). B,C) Pleural biopsies obtained by medical thoracoscopy showing necrotizing granulomas (Hematoxylin & Eosin stain). D) Acid-fast bacilli (black arrows) detected on pleural biopsies (Ziehl-Neelsen stain).

of MTB from the primary site of infection and affects predominantly immunocompromised patients, in most cases HIV-positive patients [3-5].

We hypothesize that predisposing environmental and nutritional factors related to the patient's poor living conditions contributed to a state of immunological impairment that favored mycobacterial hematogenous/lymphatic dissemination from the genital apparatus to the peritoneum and then the pleura in the case presented here, which is related to an HIV-negative patient who moved to Italy two years before [5].

Female genital tuberculosis accounts for 20-27% of cases of extra-pulmonary TB. It is a rare form of extrapulmonary TB in Europe, but a common cause of infertility in countries with a high incidence of the disease. Salpinx and endometrium are the most frequently involved organs [7,8]. During the last years, intensified migration flows from Africa to Europe have raised the number of extrapulmonary TB cases, which may be characterized, in low TB incidence countries like Italy, by unusual clinical presentations [9].

Genital TB mostly spreads from the lung or other organs by

hematogenous or lymphatic use but direct spread from near organs (*i.e.*, bowel or abdominal lymph nodes) or by sexual transmission has been described [7,8]. Notably, in our case, despite endoscopic findings being suggestive of TB, salpinx biopsies were not sent for microbiological investigations. It significantly delayed the diagnosis of MTB infection and prompt treatment start, thus enhancing mycobacterial dissemination.

Pleural tuberculosis is one of the most common forms of extrapulmonary TB worldwide [10]. It is typically paucibacillary being the genesis of pleural effusion mostly determined by a delayed hypersensitivity reaction rather than by the mycobacterial proliferation within the pleural cavity [10]. Indeed, the diagnostic performance of microbiological investigations on pleural fluid is suboptimal when compared to pleural biopsies [11,12].

Xpert MTB/RIF performed on pleural biopsies obtained by medical thoracoscopy provided rapid diagnosis and information on drug resistance, several days ahead of culture findings [13,14]. It allowed immediate treatment to start with a proper pharmacological regimen.




Figure 2. Medical thoracoscopy findings. White and confluent nodules on parietal pleura (A,B) with superior, posterolateral tent-shaped pleural adhesion (B). Diffuse white nodules on both visceral and parietal pleura (C). Thoracoscopic biopsy of a nodule of the parietal pleura (D).




Figure 3. CT scan showing pericardial (arrow) and left pleural effusion (A), and (C) parietal pleural thickening (arrow) in the left base. B) Chest X-ray performed two months after the treatment onset showing complete resolution of the pleural disease.

Medical thoracoscopy, a minimally invasive procedure performed under local anesthesia and conscious sedation, has demonstrated good accuracy in the diagnosis of tuberculous pleurisy [11,12]. It enables the selection of the most affected pleural areas for the collection of a large number of biopsies for histopathological and microbiological examinations. Furthermore, it may play an important therapeutic role in advanced pleural infections by mechanically disrupting pleural adhesions [15].

A recent study found that assessing the involvement of internal mammary lymph nodes (*via* CT scan and chest ultrasound) may be an important factor in guiding the diagnostic work-up of patients with suspected tuberculous pleurisy [16]. Notably, we did not find internal mammary lymph adenopathies; the case described here, on the other hand, is a more complex case of disseminated tuberculosis.

Conclusions

To summarize, disseminated tuberculosis is a rare and serious medical condition. Its knowledge should be implemented by healthcare workers in high-income countries. In cases of suspected disseminated TB, the diagnostic work-up should be thorough and quick to prevent MTB from spreading to multiple and/or vital organs, which could result in life-threatening events [3,4].

Microbiological examinations of biopsies of the involved organs, combined with drug-resistance information, are critical for prescribing an appropriate and timely treatment regimen.

References

1. <https://www.who.int/publications/digital/global-tuberculosis-report-2021>
2. Lewinsohn DM, Leonard MK, LoBue PA, et al. Official American Thoracic Society/Infectious Diseases Society of America/Centers for Disease Control and Prevention Clinical Practice Guidelines: Diagnosis of Tuberculosis in Adults and Children. *Clin Infect Dis* 2017;64:111-5.
3. Khan FY. Review of literature on disseminated tuberculosis with emphasis on the focused diagnostic workup. *J Family Community Med* 2019;26:83–91.
4. Mondoni M, Centola M, Viganò O, et al. Chest pain and a left parasternal soft tissue swelling in an immunocompetent refugee with disseminated tuberculosis. *Int J Infect Dis* 2020;90:116-8.
5. Suárez I, Maria Fünfer S, Jung N, et al. Severe disseminated tuberculosis in HIV-negative refugees. *Lancet Infect Dis* 2019;19:e352-9.
6. European Centre for Disease Prevention and Control, WHO

- Regional Office for Europe. Tuberculosis surveillance and monitoring in Europe 2021 – 2019 data. Available from: <https://www.ecdc.europa.eu/en/publications-data/tuberculosis-surveillance-and-monitoring-europe-2021-2019-data>
7. Chowdhury NN. Overview of tuberculosis of the female genital tract. *J Indian Med Assoc* 1996;94:345-6, 361.
 8. Wang Y, Shao R, He C, Chen L. Emerging progress on diagnosis and treatment of female genital tuberculosis. *J Int Med Res* 2021;49:3000605211014999.
 9. Dara M, Solovic I, Sotgiu G, et al. Tuberculosis care among refugees arriving in Europe: a ERS/WHO Europe Region survey of current practices. *Eur Respir J* 2016;48:808–17.
 10. Shaw JA, Irusen EM, Diacon AH, Koegelenberg CF. Pleural tuberculosis: A concise clinical review. *Clin Respir J* 2018;12:1779-86.
 11. Casalini AG, Mori PA, Majori M, et al. Pleural tuberculosis: medical thoracoscopy greatly increases the diagnostic accuracy. *ERJ Open Res* 2018;4:00046-2017.
 12. Christopher DJ, Dinakaran S, Gupta R, et al. Thoracoscopic pleural biopsy improves yield of Xpert MTB/RIF for diagnosis of pleural tuberculosis. *Respirology* 2018;23:714-7.
 13. Sehgal IS, Dhooria S, Aggarwal AN, et al. Diagnostic performance of Xpert MTB/RIF in tuberculous pleural effusion: Systematic review and meta-analysis. *J Clin Microbiol* 2016;54:1133-6.
 14. Akhter N, Sumalani KK, Chawla D, Ahmed Rizvi N. Comparison between the diagnostic accuracy of Xpert MTB/Rif assay and culture for pleural tuberculosis using tissue biopsy. *ERJ Open Res* 2019;5:00065-2019.
 15. Mondoni M, Saderi L, Trogu F, et al. Medica thoracoscopy treatment for pleural infections: a systematic review and meta-analysis. *BMC Pulm Med* 2021;21:127.
 16. Levi G, Rocchetti C, Mei F, et al. Diagnostic role of internal mammary lymph node involvement in tuberculous pleurisy: a multicenter study. *Pulmonology* 2022; S2531-0437(22)00022-8. Online ahead of print.

Non-commercial use only