

7. Coelho CC, Aquino Eda S, de Almeida DC, et al. Comparative analysis and reproducibility of the modified shuttle walk test in normal children and in children with cystic fibrosis. *J Bras Pneumol* 2007;33:168-74.
8. Lima C, de Andrade A, Campos S, et al. Six-minute walk test as a determinant of the functional capacity of children and adolescents with cystic fibrosis: A systematic review. *Respir Med* 2018;137:83-8.
9. Pereira FM, Ribeiro MÂ, Ribeiro AF, et al. Functional performance on the six-minute walk test in patients with cystic fibrosis. *J Bras Pneumol* 2011;37:735-44.
10. Hassan J, van der Net J, Helders PJ, et al. Six-minute walk test in children with chronic conditions. *Br J Sports Med* 2010;44:270-4.
11. Gulmans VA, van Veldhoven NH, de Meer K, Helders PJ. The six-minute walking test in children with cystic fibrosis: reliability and validity. *Pediatr Pulmonol* 1996;22:85-9.
12. Cunha MT, Rozov T, Caitano de Olivera R, Jardim JR. Six minutes walk test in children and adolescents with cystic fibrosis. *Pediatr Pulmonol* 2006;41:618-22.
13. Vandoni M, Correale L, Puci MV, et al. Six minute walk distance and reference values in healthy Italian children: A cross-sectional study. *PLoS One* 2018;13:e0205792.
14. Geiger R, Strasak A, Tremel B, et al. Six-minute walk test in children and adolescents. *J Pediatr* 2007;150:395-9.
15. Lammers AE, Hislop AA, Flynn Y, Haworth SG. The 6-minute walk test: normal values for children of 4–11 years of age. *Arch Dis Child* 2008;93:464-8.
16. Ulrich S, Hildenbrand FF, Treder U, et al. Reference values for the 6-minute walk test in healthy children and adolescents in Switzerland. *BMC Pulm Med* 2013;13:49.
17. Farrell PM, White TB, Ren CL, et al. Diagnosis of cystic fibrosis: Consensus guidelines from the Cystic Fibrosis Foundation. *J Pediatr* 2017;184:243.
18. ATS Committee on Proficiency Standards for Clinical Pulmonary Function Laboratories. ATS Statement. Guidelines for the Six-Minute Walk Test. *Am J Respir Crit Care Med* 2002;166:111–7. Erratum in *ATS Statement: Guidelines for the Six-Minute Walk Test. Am J Respir Crit Care Med* 2016;193:1185.
19. European Cystic Fibrosis Society. ECFS Patient Registry. Annual Data Report 2017 daata. Accessed ON: 22 December 2020. Available from: https://www.ecfs.eu/sites/default/files/general-content-images/working-groups/ecfs-patient-registry/ECFS-Report2017_v1.3.pdf
20. Quanjer PH, Stanojevic S, Cole TJ, et al. Multi-ethnic reference values for spirometry for the 3–95-yr age range: the global lung function 2012 equations the ERS Global Lung Function Initiative. *Eur Respir J* 2012;40:1324-43.
21. Flume PA, O'Sullivan BP, Robinson KA, et al. Cystic fibrosis pulmonary guidelines: chronic medications for maintenance of lung health. *Am J Respir Crit Care Med* 2007;176:957-69.
22. Gruber W, Orenstein DM, Braumann KM, Huls G. Health-related fitness and trainability in children with cystic fibrosis. *Pediatr Pulmonol* 2008;43:953-64.
23. Lesser DJ, Fleming MM, Maher CA, et al. Does the 6-min walk test correlate with the exercise stress test in children? *Pediatr Pulmonol* 2010;45:135-40.
24. Troosters T, Langer D, Vrijnsen B, et al. Skeletal muscle weakness, exercise tolerance and physical activity in adults with cystic fibrosis. *Eur Respir J* 2009;33:99-106.
25. Priesnitz CV, Rodrigues CV, Stumpf da Silva C, et al. Reference values for the 6-min walk test in healthy children aged 6-12 years. *Pediatr Pulmonol* 2009;44:1174-9.
26. Troosters T, Gossink R, Decramer M. Six minute walking distance in healthy elderly subjects. *Eur Respir J* 1999;14:270-4.
27. Chetta A, Pist G, Zanini A, et al. Six-minute walking test in cystic fibrosis adults with mild to moderate lung disease: comparison to healthy subjects. *Respir Med* 2001;95:986-91.
28. Gomez Punter RM, Girón Moreno RM, Vázquez Espinosa E, et al. Desaturations during 6-minute walk test and predicting nocturnal desaturations in adult subjects with cystic fibrosis. *Respir Care* 2019;64:48-54.
29. Rayment JH, Stanojevic S, Davis SD, et al. Lung clearance index to monitor treatment response in pulmonary exacerbations in preschool children with cystic fibrosis. *Thorax* 2018;73:451-8.
30. Saad HB, Prefaut C, Missaoui R, et al. Reference equation for 6-min walk distance in healthy North African children 6-16 years old. *Pediatr Pulmonol* 2009;44:316-24.

Non-commercial use only