

References

1. World Health Organization. Global health estimates: life expectancy and leading causes of death and disability. Available from: <https://www.who.int/data/gho/data/themes/mortality-and-global-health-estimates>.
2. Collet JP, Thiele H, Barbato E, et al. 2020 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation. *Eur Heart J* 2021;42:1289-367.
3. Morici N, De Servi S, De Luca L, et al. Management of acute coronary syndromes in older adults. *Eur Heart J* 2022;43:1542-53.
4. Tahhan AS, Vaduganathan M, Greene SJ, et al. Enrollment of older patients, women, and racial/ethnic minority groups in contemporary acute coronary syndrome clinical trials: a systematic review. *JAMA Cardiol* 2020;5:714-22.
5. van Wyk GW, Berkovsky S, Navarro DF, Coiera E. Comparing health outcomes between coronary interventions in frail patients aged 75 years or older with acute coronary syndrome: a systematic review. *Eur Geriatr Med* 2022;13:1057-69.
6. Lawton JS, Tamis-Holland JE, Bangalore S, et al. 2021 ACC/AHA/SCAI guideline for Coronary artery revascularization: a report of the American College of cardiology/American Heart Association Joint Committee on clinical practice guidelines. *J Am Coll Cardiol* 2022;79:e21-129.
7. Bamford JM, Sandercock PAG, Warlow CP, Slattery J. Interobserver agreement for the assessment of handicap in stroke patients. *Stroke* 1989;20:828.
8. Leonardi S, Bueno H, Ahrens I, et al. Optimised care of elderly patients with acute coronary syndrome. *Eur Hear J Acute Cardiovasc Care* 2018;7:287-95.
9. Tegn N, Abdelnoor M, Aaberge L, et al. Invasive versus conservative strategy in patients aged 80 years or older with non-ST-elevation myocardial infarction or unstable angina pectoris (after eighty study): an open-label randomised controlled trial. *Lancet* 2016;387:1057-65.
10. Batchelor WB, Anstrom KJ, Muhlbaier LH, et al. Contemporary outcome trends in the elderly undergoing percutaneous coronary interventions: results in 7,472 octogenarians. *J Am Coll Cardiol* 2000;36:723-30.
11. Wang TY, Gutierrez A, Peterson ED. Percutaneous coronary intervention in the elderly. *Nat Rev Cardiol* 2011;8:79-90.
12. Gimbel M, Qaderan K, Wittenstein L, et al. Clopidogrel versus ticagrelor or prasugrel in patients aged 70 years or older with non-ST-elevation acute coronary syndrome (POPular AGE): the randomised, open-label, non-inferiority trial. *Lancet* 2020;395:1374-81.

Non-commercial use only