



## Monaldi Archives for Chest Disease

eISSN 2532-5264

<https://www.monaldi-archives.org/>

**Publisher's Disclaimer.** E-publishing ahead of print is increasingly important for the rapid dissemination of science. The **Early Access** service lets users access peer-reviewed articles well before print / regular issue publication, significantly reducing the time it takes for critical findings to reach the research community.

These articles are searchable and citable by their DOI (Digital Object Identifier).

The **Monaldi Archives for Chest Disease** is, therefore, e-publishing PDF files of an early version of manuscripts that have undergone a regular peer review and have been accepted for publication, but have not been through the typesetting, pagination and proofreading processes, which may lead to differences between this version and the final one.

The final version of the manuscript will then appear in a regular issue of the journal.

E-publishing of this PDF file has been approved by the authors.

*All legal disclaimers applicable to the journal apply to this production process as well.*

Monaldi Arch Chest Dis 2023 [Online ahead of print]

*To cite this Article:*

Parato VM, Belleggia S, Parato AG, et al. **Multi-valve Libman-Sacks's endocarditis-related multiple, massive and fatal systemic embolization. A case report and a review of diagnostic work-up.** *Monaldi Arch Chest Dis* doi: 10.4081/monaldi.2023.2729

 ©The Author(s), 2023  
Licensee [PAGEPress](#), Italy

Note: The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.



# **Multi-valve Libman-Sacks's endocarditis-related multiple, massive and fatal systemic embolization. A case report and a review of diagnostic work-up**

Vito Maurizio Parato,<sup>1,2</sup> Sara Belleggia,<sup>2</sup> Andrea Giovanni Parato,<sup>3</sup> Umberto Ianni,<sup>1</sup>  
Michela Molisana,<sup>1</sup> Germana Gizzi,<sup>1</sup> Simone D'Agostino,<sup>1</sup> Melissa Dottori,<sup>4</sup>  
Marco Di Eusanio<sup>5</sup>

<sup>1</sup>Cardiology and Rehabilitation Unit, Emergency Department, Madonna del Soccorso Hospital, San Benedetto del Tronto

<sup>2</sup>Post-Graduation School of Cardiovascular Diseases, School of Medicine, Università Politecnica delle Marche, Ancona

<sup>3</sup>Cardiology Division and Chair, "Tor Vergata" University Hospital, Rome

<sup>4</sup>Cardiology Unit and Echolab, Cardiovascular Sciences Department, University Hospital Lancisi-Salesi, Ancona

<sup>5</sup>Cardiac Surgery Unit and Cardiovascular Sciences Department, University Hospital Lancisi-Salesi, Università Politecnica delle Marche, Ancona, Italy

**Corresponding Author:** Vito Maurizio Parato, MD, FESC, FACC,

Cardiology and Rehabilitation Unit, Emergency Department, Madonna del Soccorso Hospital, AST-AP, Via Luciano Manara 7, 63074 San Benedetto del Tronto (AP), Italy

Tel./Fax +39 0735793277

Email: [vitomaurizio.parato@sanita.marche.it](mailto:vitomaurizio.parato@sanita.marche.it)

## **Ethical statement**

The Ethics Committee of Ancona (Italy) waived the need for formal ethical approval since the patient's informed consent was obtained to submit this case report while maintaining anonymity.

## **Declaration of patient consent**

The authors certify that they have obtained all appropriate patient consent forms. In the form, the legal guardian has given her consent for images and other clinical information to be reported in the journal. Anonymity was guaranteed.

## **Competing interests – Funding**

No competing interests. NO FUNDING.

## **Authors' contributions**

Sara Belleggia: text writing; Andrea Giovanni Parato: references setting; Umberto Ianni: text writing; Michela Molisana: figures setting; Germana Gizzi: english form setting; Simone D'Agostino: patient's data collection; Melissa Dottori: videos setting; Vito Maurizio Parato and Marco Di Eusanio: final supervision.

## **Abstract**

We reported a case of non-bacterial thrombotic endocarditis (NBTE) in a 37-year-old woman who presented with signs and symptoms of cardio-embolic cerebral stroke caused by a prothrombotic state due to underlying advanced uterine cancer. Multimodal imaging, including 3D-ecocardiography, as well as laboratory and cultural tests, were critical in making the diagnosis. After starting anticoagulation therapy with low molecular weight heparin (LMWH), the patient underwent surgical aortic valve replacement due to worsening aortic valve function, initial left ventricle enlargement, increasing dimensions, and mobility of vegetations. Unfortunately, vegetations relapsed on the aortic valve bio-prosthesis as well as the mitral leaflets, resulting in a final picture of multi-valve NBTE. The fatal outcome was due to a massive multiple limb embolism, which resulted in leg amputations and septic complications. Starting with the case, we present a brief overview of the pathology's presentation, treatment, management, and prognosis, as well as the diagnostic work-up.

## **Introduction**

Non-bacterial thrombotic endocarditis (NBTE), also known as marantic, Libman-Sacks, thrombotic, or verrucous endocarditis, is a rare condition of non-infective endocarditis characterized by the formation of aggregates of platelets and fibrin that affects usually undamaged heart valves in the absence of bacteremia and mostly occurs in patients with a predisposing state like hypercoagulability [1-3]. NBTE is usually associated with advanced malignancy (typically adenocarcinomas), autoimmune/rheumatologic

diseases, such as systemic lupus erythematosus (SLE) and rheumatoid arthritis. Diagnosis is often post-mortem. Clinical manifestations are usually related to systemic embolization, with cerebrovascular stroke as the most common first presentation. We describe a case of NBTE in a young woman affected by advanced uterine cancer.

### **Case Report**

A 37-year-old female patient, diagnosed with advanced uterine cancer (poorly differentiated Human Papilloma Virus [HPV]-related adenocarcinoma) and recently presenting left popliteal deep vein thrombosis complicated by pulmonary embolism, presented to our Emergency Department with recurrent episodes of dysarthria and blurred vision.

Upon admission, she was referred to the Neurological Unit for comprehensive assessment. The patient denied any previous history of fever or cardiovascular symptoms. Physical examination disclosed mild dysarthria and left buccal rhyme hypokinesia.

Laboratory tests showed: low hemoglobin concentration (9.6 g/dL; normal range: 12.5 - 16.0 g/dL), elevated D-dimer levels (22.5 mg/mL; normal range <0.5 mg/dL), and an elevated C-reactive protein level (18.3 mg/dL; normal range <0.5 mg/dL). Kidney function was normal (eGFR: 90 mL/min/1.73 m<sup>2</sup>). Liver function was normal (GPT: 26 U/L, normal range: 7-45; GOT: 21 U/L, normal range: 8-43). Coagulation parameters were normal: aPTT 32 sec; INR: 1. Procalcitonin and blood cultures – for both aerobic and anaerobic bacteria - were negative. Serological tests for immunity such as rheumatoid factor, antinuclear antibody, and anticardiolipin antibody were negative.

Imaging studies, including chest and abdomen computed tomography (CT), demonstrated pulmonary embolism with an extended area of parenchymal infarction, appearance of splenic infarction, enlargement of the uterine mass with lymphnodes and caval vein invasion. Additionally, CT-brain scan showed new cortical-subcortical hypodensities indicative of recent ischemic lesions, which were confirmed by brain magnetic resonance imaging (MRI), suggesting cerebral embolism.

Due to the potential source of emboli, the patient was referred to the Cardiology Unit. Trans-thoracic echocardiography (TTE) revealed mild aortic insufficiency. In order to rule out the presence of a patent foramen ovale (PFO), TTE bubble test was performed with negative result. Therefore, a 2D-trans-esophageal echocardiography (TEE) with bubble

test was conducted, which turned out negative for PFO. However, TEE surprisingly revealed an oval-shaped isoechoic thickening of the aortic valve cusp tips, leading to mild aortic insufficiency with double jet regurgitation.

Given the patient's clinical history, laboratory results, and echocardiographic findings, there was a suspicion of marantic endocarditis. The patient was initiated on anticoagulation therapy using low molecular weight heparin (LMWH; enoxaparin, 100UI/kg bid) and also received carboplatin and paclitaxel chemotherapy for the management of advanced cancer. Bevacizumab was excluded from the treatment plan due to the thromboembolic condition. Additionally, empiric antibiotics were administered.

Two weeks later, TEE exam showed multiple isoechoic masses involving all the aortic cusps, which had increased in size. The largest mass, measuring 15 mm x 4 mm, demonstrated significant mobility with projection into the outflow tract during diastole and onto the aortic valve during systole (Figure 1 A,B; Video 1). The abnormal thickening of valve free margin caused leaflet retraction leading to significant aortic insufficiency due to coaptation gap (Figure 1 C,D; Video 2). No masses were found on the remaining valves.

Considering the poor response to medical therapy and the worsening valvular function with initial left ventricle enlargement (end-systolic diameter of 47 mm), the patient was referred to Cardiac Surgery Unit. The patient underwent aortic valve replacement with a bio-prosthesis (Carpentier-Edwards Magna Ease N. 21).

The surgical specimen was sent for histological analysis. The result was: finely granular, fibrinous eosinophilic material, containing cellular debris including nuclear remnants, neutrophils, lymphocytes.

However, two months later, she presented a severe limb ischemia, which led to the amputation of both legs. 2D-3D TEE demonstrated a relapse of new vegetations on the prosthetic aortic valve (Figure 2 A,B; Videos 3 and 4) and on both mitral leaflets (Figure 2 C,D and Videos 5 and 6). The largest mitral vegetation was 12 x 9 mm and at trans-gastric short-axis view (0°) a moderate regurgitation with central jet was visible (Video 7).

Infective endocarditis was definitely excluded because of the lack of Duke's criteria according to the latest guidelines [4].

Despite efforts to manage the situation, the patient's condition continued to deteriorate, and the patient died because of refractory sepsis one month later.

## **DISCUSSION and REVIEW**

Non-bacterial thrombotic endocarditis (NBTE), also known as *marantic*, Libman-Sacks, thrombotic or verrucous endocarditis, is a form of non-infective endocarditis characterized by the formation of aggregates of platelets plus fibrin. It usually affects undamaged heart valves without the presence of bacteremia and usually occurs in patients with a specific predisposition such as hypercoagulability [5,6].

NBTE is usually associated with advanced malignancies (80% of cases) [1,3] typically adenocarcinomas, autoimmune/rheumatic diseases such as systemic lupus erythematosus (SLE) and rheumatoid arthritis [7].

The diagnosis of NBTE can be challenging and it is often made postmortem (with rates ranging from 0.9 to 1.6%) [8-13]. The most frequent clinical presentations are related to systemic embolization of the friable vegetations, leading to conditions such as cerebrovascular transient ischemic attacks (TIAs) and strokes (the most common), myocardial ischemia, spleen infarction, hematuria, limb ischemia and ischemic skin manifestations [6,14,15].

To diagnose NBTE, it is crucial to exclude infective endocarditis (IE), considering the possibility of a culture-negative infective endocarditis. NBTE should be suspected in patients with malignancies or autoimmune diseases who develop embolic complications. Cancer screening (e.g., chest and abdomen computed tomography) is mandatory in many cases and autoimmune tests should be included into the diagnostic work-up [15,16].

Transthoracic echocardiography (TTE) and transesophageal echocardiography (TEE) represent the first-choice imaging modalities for the diagnosis of NBTE. The most common echocardiographic finding is the presence of small hyperechoic (isoechoic in our case) masses, typically attached to the low-pressure side of the heart valves, with or without oscillating motion. The mitral valve is involved in two-thirds of cases, while the aortic valve is affected in up to a quarter of cases [6,17]. Atypical presentations with larger lesions or diffuse leaflet thickening affecting multiple valves are also possible [16]. Unlike infective endocarditis, local complications such as abscesses, pseudoaneurysms, and fistulas are typically absent [18]. According to Roldan *et al.* [19] the use of 3D-TEE,

compared to 2D-TEE, provides additional information about the aortic valve more often than for the mitral valve. In fact, the number of vegetations identified and the different locations of vegetations might be higher using 3D-TEE as compared with 2D-TEE and these differences are generally most remarkable for the aortic valve (this is likely explained by the superior visualization of the bodies of the aortic cusps using 3D-TEE). Moreover, 3D-TEE detects more often associated commissural fusion of the mitral leaflets or aortic cusps.

Galzerano *et al.* [20] said that anatomic imaging and hence anatomy-based vocabulary makes three-dimensional TEE the echographic modality better able to improve the communication within the endocarditis team. Furthermore, the authors suggest performing it routinely in institutions in order to fill the gap between 2D-echocardiography and anatomy.

The management of NBTE involves anticoagulation therapy in order to reduce the risk of embolic manifestations. Heparin (unfractionated or low molecular weight) is preferred to vitamin K antagonists [6,21,22]. The routine use of direct oral anticoagulants (DOACs) for this indication has not been adequately evaluated [18] and recurrent thromboembolism after anticoagulant discontinuation has been reported, suggesting long-term continuation of anticoagulation [22,23].

Additionally, the treatment of the underlying disease (e.g., malignancy or autoimmune disorder) should be promptly initiated [6].

Surgical indications for valve intervention are not well-defined but seem to align with those of infective endocarditis, such as heart failure or acute valve rupture, and valve preservation may be possible, unlike in IE [6].

If antibiotics were initiated, they can be discontinued unless there is a suspicion of infective endocarditis with negative blood cultures.

The optimal follow-up is not well-established, but it is advisable to repeat echocardiographic evaluations to monitor the progression or resolution of valvular vegetations [24]. Prognosis is generally poor and closely related to the underlying disease, especially in cases of advanced malignancies, with a mortality rate of 38.1% [17].

## Conclusions

We've reported a case of NBTE presenting with signs and symptoms of cardio-embolic cerebral stroke and related to a prothrombotic state due to underlying advanced uterine cancer. Besides laboratory and cultural tests, multimodal imaging, including 3D-ecocardiography, was crucial for the diagnosis. Unlike other published cases [23], given the worsening aortic valve function with initial hemodynamic impact on left ventricle volumes, increasing dimensions and mobility of vegetations, after starting anticoagulation therapy, patient underwent surgical aortic valve replacement.

Unfortunately, this resulted a failing decision, since a relapse of vegetations occurred on aortic valve bio-prosthesis and, additionally, on the mitral leaflets with a final picture of multi-valve NBTE. A massive multiple limb embolism caused leg amputations and septical complications, resulting in death.

The uniqueness of our case is the multi-valve involvement and the high burden of multisite peripheral embolism caused by a significant hypercoagulable status, resulting in deep vein thrombosis and pulmonary embolism, splenic infarction, and a massively embolizing NBTE.

## References


1. Baglin T, Gray E, Greaves M, et al. Clinical guidelines for testing for heritable thrombophilia. *Br J Haematol* 2010;149:209-20.
2. Rodriguez EA, Choudhry MW, Boor PJ, et al. Primary nonbacterial thrombotic endocarditis presenting with bowel infarction secondary to superior mesenteric artery embolism. *Methodist DeBakey Cardiovasc J* 2018;14:228.
3. Venepally NR, Arsanjani R, Agasthi P, et al. A new insight into nonbacterial thrombotic endocarditis: a systematic review of cases. *Anatol J Cardiol* 2022;26:743.
4. Habib G, Lancellotti P, Antunes MJ, et al. 2015 ESC Guidelines for the management of infective endocarditis: The Task Force for the Management of Infective Endocarditis of the European Society of Cardiology (ESC). Endorsed by: European Association for Cardio-Thoracic Surgery (EACTS), the European Association of Nuclear Medicine (EANM). *Eur Heart J* 2015;36:3075-123.
5. Smeglin A, Ansari M, Skali H, et al. Marantic endocarditis and disseminated intravascular coagulation with systemic emboli in presentation of pancreatic cancer. *J Clin Oncol* 2008;26:1383-5.
6. el-Shami K, Griffiths E, Streiff M. Nonbacterial thrombotic endocarditis in cancer patients: pathogenesis, diagnosis, and treatment. *Oncologist* 2007;12:518-23.

7. Edoute Y, Haim N, Rinkevich D, et al. Cardiac valvular vegetations in cancer patients: a prospective echocardiographic study of 200 patients. *Am J Med* 1997;102:252-8.
8. Deppisch LM, Olusegun Fayemi A. Non-bacterial thrombotic endocarditis: clinicopathologic correlations. *Am Heart J* 1976;92:723-9.
9. Rosen P, Armstrong D. Nonbacterial thrombotic endocarditis in patients with malignant neoplastic diseases. *Am J Med* 1973;54:23-9.
10. Eiken PW, Edwards WD, Tazelaar HD, Zehr KJ. Surgical pathology of nonbacterial thrombotic endocarditis in 30 patients, 1985-2000. *Mayo Clin Proc* 2001;76:1204-12.
11. Llenas-García J, Guerra-Vales JM, Montes-Moreno S, et al. [Nonbacterial thrombotic endocarditis: clinicopathologic study of a necropsy series]. [Article in Spanish]. *Rev Esp Cardiol* 2007;60:493-500.
12. Steiner I. [Nonbacterial thrombotic endocarditis--a study of 171 case reports]. [Article in Czech]. *Cesk Patol* 1993;29:58-60.
13. González Quintela A, Candela MJ, et al. Non-bacterial thrombotic endocarditis in cancer patients. *Acta Cardiol* 1991;46:1-9.
14. Mazokopakis EE, Syros PK, Starakis IK. Nonbacterial thrombotic endocarditis (marantic endocarditis) in cancer patients. *Cardiovasc Hematol Disord Drug Targets* 2010;10:84-6.
15. Baglin T, Gray E, Greaves M, et al. Clinical guidelines for testing for heritable thrombophilia. *Br J Haematol* 2010;149:209-20.
16. Piccioli A, Lensing AWA, Prins MH, et al. Extensive screening for occult malignant disease in idiopathic venous thromboembolism: a prospective randomized clinical trial. *J Thromb Haemost* 2004;2:884-9.
17. Zmaili MA, Alzubi JM, Kocyigit D, et al. A contemporary 20-year Cleveland Clinic experience of nonbacterial thrombotic endocarditis: etiology, echocardiographic imaging, management, and outcomes. *Am J Med* 2021;134:361-9.
18. Silbiger JJ. The valvulopathy of non-bacterial thrombotic endocarditis. *J Heart Valve Dis* 2009;18:159-66.
19. Roldan CA, Tolstrup K, Macias L, et al. Libman-Sacks endocarditis: detection, characterization, and clinical correlates by three-dimensional transesophageal echocardiography. *J Am Soc Echocardiogr* 2015;28:770-9.
20. Galzerano D, Kinsara AJ, Di Michele S, et al. Three dimensional transesophageal echocardiography: a missing link in infective endocarditis imaging? *Int J Cardiovasc Imaging* 2020;36:403-13.
21. Lopez JA, Ross RS, Fishbein MC, Siegel RJ. Nonbacterial thrombotic endocarditis: a review. *Am Heart J* 1987;113:773-84.
22. Whitlock RP, Sun JC, Fremez SE, et al. Antithrombotic and thrombolytic therapy for valvular disease: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed:


American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. Chest 2012;141:e576S-e600S.

23. Rogers LR, Cho ES, Kempin S, Posner JB. Cerebral infarction from non-bacterial thrombotic endocarditis. Clinical and pathological study including the effects of anticoagulation. Am J Med 1987;83:746-56.
24. Yokoyama S, Iwano H, Yamada S, et al. Long-term echocardiographic evaluation of valvular lesions in a patient with nonbacterial thrombotic endocarditis associated with advanced uterine cancer. J Cardiol Cases 2016;14:82-6.

## Figures



**Figure 1.** Non-bacterial thrombotic endocarditis (NBTE) in a 37-year-old woman with uterine metastatic adenocarcinoma. **Panel A-B:** 2D-transesophageal echocardiogram, apical long axis with X-plane view, demonstrating thickening of the aortic valve leaflets together with small isoechoic masses attached to free margin (arrow). **Panel C-D:** Transesophageal echocardiogram aortic valve apical long axis and short axis view demonstrating mild aortic insufficiency with double jet regurgitation.



**Figure 2.** Non-bacterial thrombotic endocarditis (NBTE) in a 37-year-old woman with uterine metastatic adenocarcinoma. **Panel A-B:** 2D and 3D transesophageal echocardiogram (short axis view) demonstrating pathological thickening of the aortic bio-prosthetic valve leaflets together with small isoechoic masses attached to the prosthetic leaflets. **Panel C-D:** 2D (C) and 3D (D) transesophageal echocardiogram, 180° and surgical view respectively, demonstrating non-bacterial vegetations on both mitral leaflets (arrows).

---

**Video 1.** 2D-TEE X-plane view, demonstrating oval-shaped isoechoic and mobile masses on aortic valve.

**Video 2.** 2D-TEE short-axis view (60°) demonstrating aortic valve regurgitation.

**Video 3.** 2D-TEE short-axis view (60°) demonstrating new onset pathological thickening of aortic bio-prosthetic leaflets.

**Video 4.** 3D-TEE short-axis view (60°) demonstrating new onset pathological thickening of aortic bio-prosthetic leaflets.

**Video 5.** 2D-TEE X-plane view demonstrating oval-shaped isoechoic masses on both mitral leaflets.

**Video 6.** 3D-TEE surgical view demonstrating pathological thickening and masses on both mitral leaflets.

**Video 7.** 2D-TEE trans-gastric view (0°) demonstrating mitral valve involvement with significant regurgitation.