

Respiratory and physical therapy in the intensive care unit after liver transplantation for acute-on-chronic liver failure: a case report

Davide Tarello,^{1,2} Francesca Giogà,^{1,2} Andrea Lauterio,^{3,4} Chiara Becchetti,⁵ Giovanni Perricone,⁵ Giorgio Santi,⁶ Monica Ragazzi,⁷ Gianpaola Monti,⁶ Marta Lazzeri^{1,2}

¹Respiratory Physiotherapy Team, ASST Grande Ospedale Metropolitano Niguarda, Milan; ²Italian Association of Respiratory Physiotherapists (ARIR), Milan; ³Department of General Surgery and Transplantation, Niguarda Transplant Center, ASST Grande Ospedale Metropolitano Niguarda, Milan; ⁴School of Medicine and Surgery, University of Milano-Bicocca, Milan; ⁵Hepatology and Gastroenterology Unit, Niguarda Transplant Center, ASST Grande Ospedale Metropolitano Niguarda, Milan; ⁶Department of Anesthesia, Intensive Care and Transplantation, Niguarda Transplant Center, ASST Grande Ospedale Metropolitano Niguarda, Milan; ⁷Multi-specialist Medical Department, ASST Grande Ospedale Metropolitano Niguarda, Milan, Italy

Correspondence: Davide Tarello, Respiratory Physiotherapy Team, ASST Grande Ospedale Metropolitano Niguarda, Piazza Ospedale Maggiore 3, 20162 Milan, Italy.
Tel.: +39 347 2752893.
E-mail: tarello davide.fisioterapista@gmail.com

Key words: case report, liver transplantation, acute-on-chronic liver failure, intensive care unit, physiotherapy, extubation failure, non-invasive ventilation.

Contributions: TD, GF, were the physiotherapists in charge of the patient, data collection and writing the manuscript. LA, BC, PG, RM, were the physicians in charge of the patient, data collection, and reviewing the paper; SG, was the nurse who took care of the patient, collected data, and reviewed the paper; LM, MG, supervised the project. All authors contributed to the final manuscript.

Conflict of interest: all authors declare there is no conflict of interest.

Ethics approval and consent to participate: no ethical committee approval was required for this case report by the Department because this article does not contain any studies with human participants or animals. Informed consent was obtained from the patient included in this study.

Patient consent for publication: the patient gave her written consent to use his personal data for the publication of this case report and any accompanying images.

Availability of data and materials: all data underlying the findings are fully available.

Funding: none.

Acknowledgments: the authors wish to thank all the ICU staff (MDs, nurses, and physiotherapists) for their excellent job and cooperation in the care of this patient. A special acknowledgment to the patient, who immediately gave his permission to write and publish this paper.

Received: 2 January 2024.

Accepted: 20 July 2024.

Early view: 10 September 2024.

Publisher's note: all claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

©Copyright: the Author(s), 2024

Licensee PAGEPress, Italy

Monaldi Archives for Chest Disease 2025; 95:2898

doi: 10.4081/monaldi.2024.2898

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial International License (CC BY-NC 4.0) which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

Abstract

Acute-on-chronic liver failure (ACLF) is a severe clinical condition for which liver transplantation (LT) is the only curative option. Due to the recipients' generally poor pre-operative clinical conditions and extensive surgery, post-LT respiratory disorders are very common and significantly contribute to related morbidity and mortality.

We report the case of a 49-year-old patient with ACLF grade 3 who has been taken care of by the Respiratory Physiotherapy Team since hospital admission. After the extubation, the patient was supported with non-invasive ventilation and mechanical in-exsufflation; meanwhile, early resistance and functional training were started. No adverse events occurred during physiotherapy sessions, and the patient returned home without respiratory support.

Respiratory and physical therapy in the intensive care unit after LT were safe and feasible interventions for this patient. Given the high incidence of postoperative pulmonary complications and the high rehabilitation needs, we suggest that physiotherapy should be provided for ACLF recipients.

Introduction

Acute-on-chronic liver failure (ACLF) is a clinical syndrome defined by an acute deterioration of liver function associated with extrahepatic organ failures often requiring intensive care support and associated with a high rate of short-term mortality [1]. The global prevalence of ACLF among patients admitted with decompensated cirrhosis is estimated at 35% [2]. Liver transplantation (LT) is considered the only life-saving treatment that can potentially change the long-term prognosis of patients with ACLF [3]. Due to recipients' generally poor pre-operative clinical conditions and to the extensive surgery, post-LT respiratory disorders are very common and significantly contribute to the related morbidity and mortality, with lung failure being itself an independent predictor of in-hospital mortality in ACLF patients [4,5]. It is also acknowledged that healthcare resource utilization after LT was greater among

recipients with ACLF compared with patients without ACLF. They experienced longer median post-LT length of hospital stay (9 vs. 39 days), longer intensive care unit (ICU) stay (5 vs. 11 days), and a higher rate of discharge to a rehabilitation center (16% vs. 30%) [6,7]. Currently, despite the relevance of the topic, no evidence exists about the efficacy and modalities of rehabilitation intervention in ACLF patients receiving transplantation. We present the case of a 49-year-old patient with ACLF grade 3 admitted to our LT center for evaluation, who was taken care of by the respiratory physiotherapy team from day 1 until his discharge from the hospital.

The aim of this paper, which follows the Case Report (CARE) Guidelines [8], is to describe the respiratory and physical therapy interventions in the ICU setting after LT in this ACLF patient.

Case Report

We report the case of a 49-year-old patient with ACLF grade 3 who was admitted to our ICU for evaluation for LT after 26 days of hospitalization (of which 14 were in the ICU) (Figure 1). At admission, the patient had an ACLF grade 3 with four organ failures (brain, circulation, liver, and kidney), a Chronic-Liver Failure C score of 73 (30-day mortality 90%, 3-month mortality 97%), and a model for End-Stage Liver Disease score of 40. The past clinical history was mute, except for a former ultrasound documenting liver steatosis. The patient was an active smoker and active drinker. Harmful alcohol consumption was identified as a precipitant of ACLF.

As soon as the patient arrived in our transplantation center (where physiotherapy referral is automatic), the physiotherapist on duty instantly placed the patient's limbs in neutral position with anti-decubitus devices and verified the patient's bed position to prevent the worsening of pre-existing pressure injuries and neuromuscular disorders. In the pre-transplantation phase, preventive respiratory physiotherapy interventions were provided, including continuous active humidification (at 37°C) of the airways, changing

patient's decubitus (on the lateral decubitus and "in-bed" sitting position) and airway clearance techniques (such as lung expansion and recruitment techniques and expiratory rib cage compressions) to prevent mucus encumbrance and atelectasis and to improve gas exchange.

After surgery, the patient's unstable clinical conditions (hepatic encephalopathy, kidney failure, need for vasoactive drugs) led to a slow-paced weaning from mechanical ventilation and cardiovascular support. From day 0 to post-operative day (POD) 5, the physiotherapist regularly evaluated the patient's neuromotor and respiratory functions and, eventually, his readiness to manage spontaneous breathing. Firstly, to evaluate the neurologic status, Richmond Agitation Sedation Scale, Confusion Assessment Method for the ICU, and ability to follow simple orders were considered, while motor function was assessed by using the Medical Research Council Sum Score (MRC-SS). Secondly, to evaluate the patient's respiratory function, tidal volume, respiratory rate, respiratory kinematics, P0.1, Pmusc index, Rapid Shallow Breathing Index, arterial blood gas, presence of voluntary cough and cough reflex, bronchial secretions quality and quantity, and negative inspiratory force (NIF), where considered, and a Spontaneous Breathing Trial (SBT) was performed. On POD 6, after successfully passing the SBT and the cuff-leak test, the patient was extubated. Since the high risk of extubation failure due to the presence of copious bronchial secretions, pleural effusion, and atelectasis, and to the severe muscle weakness (clinical picture of severe ICU-acquired weakness with MRC-SS 34/60 and inspiratory muscles and cough weakness with NIF-22 cmH₂O), the patient was supported with Non-invasive ventilation (NIV), high-flow nasal cannula (HFNC), and mechanical in-sufflation (MI-E). Non-invasive respiratory supports (NIRS) were primarily managed by the physiotherapist, who was responsible for the initial settings (along with the physician), choice of the most appropriate NIV mask and ventilator, and monitoring of patient response (along with the nurse). MI-E was set and firstly administered by the physiotherapist, too, then on demand by physiotherapists or nurses. To do all these tasks and to monitor patient




Figure 1. Visual representation of the hospitalization of the patient. Time 0 is considered the day of surgery. The area under the curve visually represents the rehabilitation and assistance needs in terms of respiratory (respiratory therapy - blue) and neuro-motor (physical therapy - yellow) interventions. ICU, intensive care unit; HDU, high dependency unit; POD, post-operative day.

response, the physiotherapist was present all-day shift (ICU dedicated) and 10 hours on-call. On POD 8, patients suffered from a massive secretion encumbrance causing respiratory distress, nonresponsive to MI-E (probably due to the high respiratory rate and patient-device asynchrony) or tracheal aspirations (which unfortunately led to upper airway bleeding); therefore, the patient needed to be reintubated. On POD 12, the second extubation attempt took place: even this time, the patient was supported with NIV (especially during night rest), HFNC during NIV breaks, and MI-E. On POD 13, swallowing function was assessed for screening using the Gugging Swallowing Screen for the ICU (GUSS-ICU) test which tested positive; therefore, a speech and language therapy consult was requested for formal evaluation and rehabilitation. The feeding tube was kept in place as a parallel nutritional support until the patient reacquired a proper swallowing ability. On the same day, resistance and functional training were started (initially with an in-bed cycle ergometer) (Figure 2), while functional outcome, as sitting in a wheelchair, was reached on POD 14. On POD 22, the patient was transferred to the High Dependency Unit, where the Respiratory and Physical Therapy program was carried out by the same physiotherapy team (NIV at night, MI-E for airway clearance, continuation of exercise training).

No adverse events occurred during physiotherapy sessions, which continued after the discharge from the hospital (POD 97). Finally, the patient returned home without respiratory support, walking with assistive devices.

Discussion

This case report fully describes the physiotherapy care of a frail ACLF recipient in the ICU setting.

One of the main goals of the respiratory physiotherapist in this setting is to prevent and treat respiratory complications and lung failure, which is known to be an independent predictor of in-hospital mortality in ACLF recipients [5]. In critically ill subjects, the worsening of the respiratory function can often be related to pulmonary secretion retention and/or atelectasis. In fact, in the mechanically ventilated patient, due to the presence of the artificial airway, poor humidification of inspired gases, and relative immobility, the primary mechanisms of secretion clearance are impaired [9]. Since adequate conditioning of inhaled gases can improve mucociliary clearance function and mucus rheology, active humidification was adopted as the first-line treatment for obstructive atelectasis preven-



Figure 2. Physiotherapy session in the intensive care unit: the patient is supported with non-invasive ventilation during in-bed cycle ergometry training.

tion. Expiratory Ribe Cage Compressions were also used as an add-on airway clearance technique on account of lung auscultation and ventilatory waveforms analysis [9]. Decubitus changing and positional therapy, which are essential to prevent the formation of bedsores or promote their treatment when already present, in this specific case, were implemented primarily as an adjuvant in strategies for airway clearance and lung recruitment [9]. It is well known that the recumbent position in critical patients affects ventilation, causing decreased lung volume and increased airway resistance, contributing to hypoxemia; early mobilization, instead, reduces the negative effects associated with bed rest [10,11]. In fact, it has been observed that in critically ill subjects, the sitting position leads to an increase in functional residual capacity and an improvement in oxygenation [10]. Early mobilization is a safe intervention that can decrease the incidence of intensive care unit acquired weakness (ICU-AW), increase the number of ventilator-free days, and the discharged-to-home rate for patients with a critical illness [11]. In this population, early mobilization and rehabilitation are strongly recommended by the Enhanced Recovery After Surgery (ERAS) guidelines, which state that they should be encouraged with early-goal-directed interventions, from the morning after LT until hospital discharge [12]. Unfortunately, in this specific case, due to the recipient's poor neurological and respiratory function, it was possible to start active mobilization only on POD 13 (27 days after the first ICU admission), when he had already developed an acquired weakness. Despite this, it can be said that the rehabilitation intervention was successful, as the patient returned home without respiratory support, walking independently with an assistive device.

Along with exercise therapy, an accurate nutritional intervention was carried out; the feeding tube was kept in place as a parallel nutritional support until the patient re-acquired a proper swallowing ability. This aspect, too, was fundamental for the patient's rehabilitation.

This case report also highlights the central role physiotherapists played in the respiratory weaning of this frail patient. In our hospital, we have a specialized team of respiratory and critical care physiotherapists, who, after attending a university master's course and/or post-graduate courses, have an extended knowledge and scope of practice in the cardiorespiratory, surgical, and critical care field. They have expertise to assess cough strength, work of breathing, respiratory muscle strength, and respiratory secretion load, which are important factors in the outcome of extubation [13]. Cork *et al.* showed that specialist physiotherapists can detect patients at high risk of extubation failure with a high sensitivity following a thorough assessment [13]. On top of that, Thille *et al.* observed that ICU-AW is associated with ineffective cough and extubation failure [14]. Accurate prediction of extubation outcome and risk stratification can help to inform management plans pre- and post-extubation and therefore reduce the likelihood of extubation failure. Based upon patient assessment, which highlighted severe muscle weakness (MRC-SS 34/60), inspiratory muscles and cough weakness (NIF -22 cmH₂O), prolonged intubation and presence of copious bronchial secretions, pleural effusion and atelectasis, extubation was classified as high risk. According to the most recent international clinical practice guidelines, which recommend the use of NIV immediately after extubation to prevent respiratory failure in patients at high risk of reintubation [15], and the use of HFNC, especially during NIV breaks [16], this patient was supported with NIRS strategies, which were managed by an experienced team composed by critical care physicians, critical care nurses and respiratory and critical care physiotherapist. As mentioned above, ineffective cough and inability to clear copious secretions are reasons for extu-

bation and NIV failure. Therefore, the adopted weaning strategy was the bundle of NIV and airway clearance techniques such as mechanical in-exsufflation, as the latter was essential for the former's effectiveness [17,18].

Post-extubation dysphagia is a condition that is becoming a growing concern. This occurs in 3-62% of extubated patients and can be related to mixed etiologies, such as neuromuscular impairment, critical illness, or laryngeal damage. Recent studies recommend the implementation of a standardized swallowing screening to prevent aspiration and decrease pneumonia rate and mortality [19]. In this case, the physiotherapists, along with the nurse in charge, assessed the patient's swallowing function using the GUSS-ICU screening tool, which tested positive; this was crucial for aspiration pneumonia prevention and led to an early request for formal speech and language therapy evaluation and related interventions.

Lastly, since no adverse events occurred during physiotherapy sessions, it can be said that respiratory and physical therapy after LT in the ICU setting were safe and feasible interventions in this ACLF recipient. Limitations of our experience are mainly related to the inability to draw robust conclusions exportable to the whole population, given the fact that we report experience with a single patient. Further studies are needed to show the safety and feasibility of early rehabilitation in larger cohorts. Lastly, our experience relates to a setting where there is high resource availability, with 10-hour on-call availability of respiratory physiotherapists, which is not necessarily applicable to all ICU settings.

Conclusions

Respiratory and physical therapy after LT in the ICU setting were safe and feasible interventions in this ACLF patient and were important components of the patient's care. Since the high incidence of post-operative pulmonary complications and the high rehabilitation needs, we suggest that a multidisciplinary approach, including physiotherapy, should be provided to ACLF recipients, especially for more advanced ACLF grades and in frail patients.

References

1. Perricone G, Artzner T, De Martin E, et al. Intensive care management of acute-on-chronic liver failure. *Int Care Med* 2023; 49:903-21.
2. Mezzano G, Juanola A, Cardenas A, et al. Global burden of disease: acute-on-chronic liver failure, a systematic review and meta-analysis. *Gut* 2022;71:148-55.
3. Belli LS, Duvoux C, Artzner T, et al. Liver transplantation for patients with acute-on-chronic liver failure (ACLF) in Europe: results of the ELITA/EF-CLIF collaborative study (ECLIS). *J Hepatol* 2021;75:610-22.
4. Feltracco P, Carollo C, Barbieri S, et al. Early respiratory complications after liver transplantation. *World J Gastroenterol* 2013;19:9271-81.
5. Laici C, Guizzardi C, Morelli MC, et al. Predictive factors of inhospital mortality for ICU patients with acute-on-chronic liver failure undergoing liver transplantation. *Eur J Gastroenterol Hepatol* 2022;34:967-74.
6. Chen GH, Wu RL, Huang F, et al. Liver transplantation in acute-on-chronic liver failure: excellent outcome and difficult posttransplant course. *Front Surg* 2022;9:914611.
7. Sundaram V, Lindenmeyer CC, Shetty K, et al. Patients with

- acute-on-chronic liver failure have greater healthcare resource utilization after liver transplantation. *Clin Gastroenterol Hepatol* 2023;21:704-12.e3.
8. Riley DS, Barber MS, Kienle GS, et al. CARE guidelines for case reports: explanation and elaboration document. *J Clin Epidemiol* 2017;89:218-35.
 9. Volpe MS, Guimarães FS, Morais CC. Airway clearance techniques for mechanically ventilated patients: insights for optimization. *Respir Care* 2020;65:1174-88.
 10. Hickmann CE, Montecinos-Munoz NR, Castanares-Zapatero D, et al. Acute effects of sitting out of bed and exercise on lung aeration and oxygenation in critically ill subjects. *Respir Care* 2021;66:253-62.
 11. Zhang L, Hu W, Cai Z, et al. Early mobilization of critically ill patients in the intensive care unit: A systematic review and meta-analysis. *PLoS One* 2019;14:e0223185.
 12. Brustia R, Monsel A, Skurzak S, et al. Guidelines for perioperative care for liver transplantation: enhanced recovery after surgery (ERAS) recommendations. *Transplantation* 2022;106:552-61.
 13. Cork G, Camporota L, Osman L, Shannon H. Physiotherapist prediction of extubation outcome in the adult intensive care unit. *Physiother Res Int* 2019;24:e1793.
 14. Thille AW, Boissier F, Muller M, et al. Role of ICU-acquired weakness on extubation outcome among patients at high risk of reintubation. *Crit Care* 2020;24:86.
 15. Rochwerg B, Brochard L, Elliott MW, et al. Official ERS/ATS clinical practice guidelines: noninvasive ventilation for acute respiratory failure. *Eur Respir J* 2017;50:1602426.
 16. Oczkowski S, Ergon B, Bos L, et al. ERS clinical practice guidelines: high-flow nasal cannula in acute respiratory failure. *Eur Respir J* 2022;59:2101574.
 17. Bach JR, Gonçalves MR, Hamdani I, Winck JC. Extubation of patients with neuromuscular weakness: a new management paradigm. *Chest* 2010;137:1033-9.
 18. Chatwin M, Toussaint M, Gonçalves MR, et al. Airway clearance techniques in neuromuscular disorders: a state of the art review. *Respir Med* 2018;136:98-110.
 19. Christensen M, Trapl M. Development of a modified swallowing screening tool to manage post-extubation dysphagia. *Nurs Crit Care* 2018;23:102-7.