

Prediction of spirometry outcome in Croatian patients with chronic obstructive pulmonary disease

Erim Bešić,¹ Davorka Muršić,² Tajana Jalušić Glunčić,² Jelena Ostojić,² Sanda Škrinjarić-Cincar,³ Martina Dokoza,⁴ Nataša Karamarković Lazarušić,⁵ Miroslav Samaržija,^{2,6} Andrea Vukić Dugac^{2,6}

¹Faculty of Pharmacy and Biochemistry, University of Zagreb; ²Clinic for Respiratory Diseases, University Hospital Center Zagreb; ³Faculty of Medicine, University of Osijek; ⁴Department of Internal Medicine, Zadar General Hospital; ⁵Polyclinic for the Respiratory Tract Diseases, Zagreb; ⁶School of Medicine, University of Zagreb, Croatia

Abstract

The current study offers an extensive examination of the influence of 29 diverse parameters on spirometry measurement variables in a cohort of 534 patients with chronic obstructive pulmonary disease (COPD) from five different centers in Croatia. The study elucidates both the magnitude and direction of the effect exerted by the 29 predictors on forced vital capacity (FVC), forced expiratory volume in one second (FEV1), the ratio FEV1/FVC, and predicted forced expiratory flow at 50% of FVC. Additionally, the development of prediction models for these parameters has been undertaken using several statistical methods.

The study identifies fat-free mass index, 6-minute walk distance, predicted diffusing capacity of the lung for carbon monoxide, arterial partial pressure of oxygen, and both arterial and tissue hemoglobin oxygen saturation percentage as robust positive predictors for all four spirometry parameters. Body mass index is recognized as a weak positive predictor for FEV1 and FEV1/FVC, commonly observed in COPD patients. As expected, smoking years is identified as a strong negative predictor for all four spirometry parameters, while age and illness duration exhibit strong predictive negative associations. Furthermore, the modified Medical Research Council, arterial partial pressure carbon dioxide, St. George's Respiratory Questionnaire, COPD assessment test, Depression Anxiety Stress Scales, and Nutritional Risk Screening are identified as weak negative predictors. The Charlson Comorbidity Index, phase angle, and number of comorbidities do not exhibit a significant impact on spirometry variables.

Ultimately, the performed factorial analysis categorized the 29 parameters into five groups, which were identified as relating to lung function, health status, nutritional status, age, and smoking. Multiple regression analysis, including four newly derived parameters based on the results of factorial analysis, identified nutritional status as a positive predictor for spirometry readings, while smoking, poor health status, and age were identified as negative predictors in successive order.

Key words: COPD, spirometry parameters, general linear model, multiple linear regression, stepwise regression, factorial analysis.

Correspondence to: Andrea Vukić Dugac, Clinic for Respiratory Diseases, University Hospital Center Zagreb, Jordanovac 104, 10 000 Zagreb, Croatia.
E-mail: adugac71@gmail.com

Introduction

Chronic obstructive pulmonary disease (COPD) is a prevalent lung ailment characterized by constrained airflow and respiratory difficulties. COPD is a heterogeneous lung condition characterized by chronic respiratory symptoms (dyspnea, cough, sputum production, and/or exacerbations) due to abnormalities of the airways (bronchitis, bronchiolitis) and/or alveoli (emphysema) that cause persistent, often progressive, airflow obstruction. It stands as one of the top three leading causes of mortality worldwide, imposing a substantial burden on global healthcare systems and public health. The progressive nature of COPD, coupled with its significant impact on quality of life and mortality rates, underscores the critical need for effective prevention, diagnosis, and management strategies [1-4].

Spirometry test variables are crucial in diagnosing and manag-

ing respiratory conditions, particularly COPD and asthma. Key variables include forced vital capacity (FVC), which measures the maximum amount of air a person can forcibly exhale after a full inhalation, and forced expiratory volume in one second (FEV1), which measures the amount of air exhaled in the first second of the FVC manoeuvre. The ratio FEV1/FVC helps determine the presence and severity of airway obstruction. These spirometry variables are essential for assessing lung function, monitoring disease progression, evaluating treatment efficacy, and guiding clinical decision-making [5-8].

An overview of studies examining the relationships between lung function and various contributing factors has revealed a complex structure underlying the conditioning of FEV1, FVC, and FEV1/FVC [9-11]. The multifaceted nature of factors influencing lung function highlights the challenges in determining accurate and reliable predicted values for FEV1, FVC, and FEV1/FVC



[12]. Most equations used to estimate predicted values of spirometry parameters typically incorporate a limited number of predictors [13]. However, it is evident that expanding the number of predictors in predictive models for spirometry parameters enhances their robustness and accuracy in clinical applications. Incorporating a broader array of predictors allows for a more comprehensive evaluation of lung function, considering both clinical and physiological variables [14].

Several studies have advocated for the inclusion of additional predictors beyond conventional factors such as age, sex, and height, to improve the accuracy of spirometry prediction equations [5,10,15,16]. For instance, factors like smoking history, comorbidities, body composition metrics, and respiratory symptoms have been shown to contribute significantly to lung function variability [17,18]. By incorporating these factors into predictive models, clinicians can obtain more personalized and precise estimates of spirometry parameters for individual patients [19].

Recently, a multicenter study was conducted across four different regions of Croatia, where COPD patients were recruited and evaluated regarding 29 various parameters. The study included 534 patients, of whom 325 (60.9%) were men and 209 (39.1%) were women, with a mean age of 66.7 ± 8.3 years. The mean duration of the disease was 6.9 ± 1.2 years. Most patients exhibited a moderate to severely abnormal obstructive pattern with a reduced 6-minute walk distance (6MWD) (396.5 ± 110.8 m). The patients were predominantly overweight or obese, with an average body mass index (BMI) of 26.4 ± 5.5 kg/m², and demonstrated satisfactory values for nutritional status variables, including fat-free mass index (FFMI), skeletal muscle mass index (SMMI), and phase angle (PhA). Primarily, the study was designed to assess the nutritional status of COPD patients and to investigate the association between nutritional status, disease severity, and exercise capacity in four different regions of Croatia [20]. The subsequent publication delves into a more targeted investigation, specifically examining obese patients diagnosed with COPD, investigating aspects related to body composition, pulmonary function tests, exercise capacity, and quality of life in these patients [21]. However, within these two publications stemming from the same cohort of COPD patients, we undertook preliminary investigations to ascertain which among the 29 measured parameters assessed exerted an influence on spirometry outcomes.

Therefore, the present study aims to elucidate the factors impacting variables derived from pulmonary function tests and to determine both the magnitude and direction (whether positive or negative) of the effects exerted by numerous predictors on each spirometry parameter. To achieve this objective, several statistical analyses were employed, including general linear models, stepwise regression models, and multiple regression models guided by the findings of factorial analysis.

Materials and Methods

Patients

This multicenter study involved a cohort of 534 patients diagnosed with COPD, who were recruited and assessed in the years 2018 and 2019. The study encompassed five distinct centers, comprising two from the Mediterranean region and three from continental Croatia. Patients with COPD were screened for eligibility and recruited during ambulatory visits at outpatient

clinics. Consecutive patients, predominantly current or former smokers aged 45 to 90 years, with a confirmed COPD in accordance with the Global Initiative for Chronic Obstructive Lung Disease (GOLD) guidelines, were included. All COPD patients enrolled in the study were required to be in a stable phase of the disease, defined as having no exacerbations for at least the preceding month, with no changes in respiratory medication and no symptoms indicative of a lower respiratory tract infection. Additional exclusion criteria included recent exacerbations, active malignant diseases, uncontrolled arterial hypertension (systolic pressure >180 mmHg, diastolic pressure >100 mmHg), presence of a pacemaker, or finger amputation. Ethical approval for the study protocol was obtained from the respective Ethics Committees, and all participating individuals provided informed consent. Diagnosis of COPD was established in accordance with the 2019 GOLD guidelines. Patients exhibiting indications of asthma or positive bronchodilator responses were excluded from the study, as were those with musculoskeletal, neurological, or other conditions that might influence the results of the 6MWD test.

For each participant, data encompassing 29 distinct parameters were meticulously collected. In addition to recording the age, sex, and disease duration of the patients, the evaluation included a comprehensive suite of assessments: pulmonary function tests, the 6-minute walking test, BMI, and body composition *via* bioelectrical impedance analysis, which provided measurements such as FFMI, SMMI, and PhA. Oxygenation status was assessed through arterial hemoglobin oxygen saturation percentage [a-Sa(Hb)-O₂%] and tissue hemoglobin oxygen saturation percentage [t-Sa(Hb)-O₂%], both measured using pulse oximetry.

Furthermore, the study captured data on smoking habits (including smoking status, smoking years, and pack-years), parameters reflecting gas exchange efficiency in the lungs — such as the predicted diffusing capacity for carbon monoxide (DLCO%), the predicted carbon monoxide transfer coefficient (KCO%), and the partial arterial pressures (PO₂) and carbon dioxide (PCO₂) — as well as measures of oxygenation status: a-Sa(Hb)-O₂% and t-Sa(Hb)-O₂%.

Additional assessments included various validated questionnaires and indices to gauge psychological, respiratory, nutritional status, and quality of life. These included the Depression Anxiety Stress Scales (DASS), St. George's Respiratory Questionnaire (SGRQ-C), Nutritional Risk Screening (NRS-2002), the BMI/air-flow obstruction/dyspnea/exercise capacity index (BODE), Charlson Comorbidity Index (CCI), GOLD classification [GOLD (I-IV)], the COPD Assessment Test (CAT), and the modified Medical Research Council dyspnea scale (mMRC).

Methods

Spirometry and the measurement of lung DLCO were conducted in accordance with the standards set by the American Thoracic Society/European Respiratory Society (ATS/ERS). The 6-minute walk test was administered following the guidelines provided in the ATS/ERS statement.

Bioelectrical impedance analysis is a widely accepted method for assessing body composition in patients with COPD. In this study, body composition measurements were obtained using the TANITA MC-780MA P device (Tanita Corporation, Tokyo, Japan), which employs an eight-contact electrode system. The procedures were performed according to the manufacturer's instructions and the recommendations outlined by Kyle *et al.* [22].



Statistical analysis

Statistical analysis was performed using Minitab 17 statisticam software (Minitab, LLC, State College, Pennsylvania, USA). For the evaluation of the effect of various predictors on each of the four dependent spirometry variables, a general linear model with three predictors (factors: predictor, sex, age) was used. Modeling the linear regression model for each of the dependent spirometry variables, which included a reduced number of predictors, was performed using the stepwise method. To condense 29 variables into a smaller number of factors, factor analysis followed by varimax rotation was employed. Multiple linear regression was utilized to create a model for spirometry variables, incorporating four of the five newly derived factors resulting from the factorial analysis. P-values less than 0.05 were considered statistically significant in all statistical methods used in this study. All computer outcomes are provided in the *Supplementary Material*.

Results and Discussion

All measured values were numerical, with most exhibiting a symmetrical or normal distribution. To determine the normality of the given dataset, the Shapiro-Wilk test was employed ($\alpha=0.05$). All non-normally distributed variables (Shapiro-Wilk: $p>0.001$) were log-transformed before inclusion in multivariable models to approximate a normal distribution. The distributions for the parameters NRS-2002, BODE, DASS, and SGRQ-C significantly deviated from normality (Shapiro-Wilk: $p<0.001$), yet they were still included in the parametric linear regression analysis alongside other parameters. Additionally, outcomes for GOLD (I-IV) were converted to numerical data (GOLD I = 1, GOLD II = 2, GOLD III = 3, GOLD IV = 4) and treated as discrete numerical variables. The same treatment was applied to smoking status (non-smokers = 1, ex-smokers = 2, smokers = 3). Accordingly, caution should be exercised when interpreting results obtained for these six predictors.

Descriptive statistics for all 29 parameters are displayed in Table 1. Normally distributed variables are reported as the mean \pm standard error, whereas non-normally distributed data are presented as the median and interquartile range. Results for gender-sensitive parameters are listed separately for males and females.

Initially, we investigated the impact of each individual parameter (referred to as independent variables, predictors, or factors) on the four spirometry variables (FVC, FEV1, FEV1/FVC, and FEF50%). Opting for a simple general linear model proved to be the most appropriate method for modeling a single independent quantitative variable with multiple quantitative dependent variables (the predictor itself, age, and sex). Moreover, utilizing the data in their original, non-standardized form would hinder direct comparisons across different independent variables due to disparities in their scales. Additionally, such variables would not facilitate comparative assessments of the effects of one independent variable in relation to another within the model. Consequently, the general linear model was conducted using standardized variables. This approach yielded standardized regression coefficients (β), which denote the alteration in the dependent variable associated with a one-unit change in the corresponding independent variable, while maintaining other variables constant. These coefficients quantitatively capture the influence of each independent variable, delineating both its direction and magnitude.

The results of the general linear model for FVC, FEV1, FEV1/FVC, and FEF50% are presented in *Tables S1-S4* in the *Supplementary Material*, respectively, accompanied by Figure 1,

Table 1. Descriptive statistics of measured variables.

Variable	
Age (years)	66.65 \pm 0.36
BMI (kg/m ²)	26.36 \pm 0.24
Diagnosis (years)	5.00 (3.00)
Smoking (years)	35.00 (10.00)
Pack-year	45.00 (17.50)
FVC (L) - total	2.81 \pm 0.04
FVC (L) - men	3.14 \pm 0.05
FVC (L) - women	2.30 \pm 0.05
FEV1 (L) - total	1.45 \pm 0.03
FEV1 (L) - men	1.60 \pm 0.04
FEV1 (L) - women	1.22 \pm 0.04
FEV1/FVC - total	0.52 \pm 0.01
FEV1/FVC - men	0.50 \pm 0.01
FEV1/FVC - women	0.54 \pm 0.01
FEF50% - total	23.31 \pm 0.78
FEF50% - men	22.47 \pm 0.84
FEF50% - women	24.63 \pm 1.51
DLCO%	57.56 \pm 0.90
KCO%	68.99 \pm 1.00
t-Sa(Hb)-O ₂ %	94.52 \pm 0.12
a-Sa(Hb)-O ₂ %	94.07 \pm 0.11
PO ₂ (mmHg)	72.57 \pm 0.45
PCO ₂ (mmHg)	39.49 \pm 0.25
6MWD (m) - total	396.53 \pm 4.80
6MWD (m) - men	397.07 \pm 6.18
6MWD (m) - women	395.70 \pm 7.65
Number of comorbidities	2.00 (1.00)
FFMI - total	19.88 \pm 0.14
FFMI - men	20.60 \pm 0.16
FFMI - women	18.75 \pm 0.27
PhA - total	6.00 (1.10)
PhA - men	6.00 (1.00)
PhA - women	5.75 (1.50)
SMMI	7.73 \pm 0.06
SMMI - men	8.23 \pm 0.08
SMMI - women	6.94 \pm 0.07
GOLD (I-IV)	2.00 (1.00)
mMRC	2.00 (1.00)
CAT	15.00 (10.00)
DASS	10.00 (14.00)
SGRQ-C	39.12 (31.00)
NRS-2002	1.00 (2.00)
BODE	3.00 (3.00)
CCI	4.00 (1.00)

BMI, body mass index; diagnosis, number of years since first diagnose; FVC, forced vital capacity; FEV1, forced expiratory volume in one second; FEV1/FVC, FEV1/FVC ratio; FEF50%, forced expiratory flow at 50% of FVC; DLCO%, predicted diffusing capacity for carbon monoxide; KCO%, predicted carbon monoxide transfer coefficient; t-Sa(Hb)-O₂%, tissue hemoglobin oxygen saturation percentage; a-Sa(Hb)-O₂%, arterial hemoglobin oxygen saturation percentage; PO₂, arterial partial pressure of oxygen; PCO₂, arterial partial pressure carbon dioxide; 6MWD, 6-minute walk distance; FFMI, adjusted fat-free mass index; PhA, phase angle; SMMI, skeletal muscle mass index; GOLD (I-IV), global initiative for chronic obstructive lung disease; mMRC, modified medical research council; CAT, chronic obstructive pulmonary disease assessment test; DASS, depression anxiety stress scales; SGRQ-C, St. George's respiratory questionnaire; NRS-2002, nutritional risk screening; BODE, body mass index/airflow obstruction/dyspnea/exercise capacity index; CCI, Charlson comorbidity index.



as well as *Figures S1-S4* provided in the *Supplementary Material*. Positive standardized β coefficients are highlighted in blue in *Tables S1-S4*, while negative β coefficients are marked in red. Predictors with non-statistically significant β coefficients are indicated in black. For more detailed information on the general linear model conducted, one can refer to the program outputs provided in the *Supplementary Material*.

The systematic presentation of all quantitative data presented in *Tables S1-S4* has been qualitatively organized in Table 2. Table 2 categorizes the predictors for all four spirometry parameters into strong, moderate, and weak (both positive and negative) categories. Strong predictors are defined as those with an absolute value of the β coefficient greater than 0.5 ($|\beta| > 0.5$), moderate predictors are those with a β coefficient value between 0.3 and 0.5 in absolute terms ($0.3 \leq |\beta| \leq 0.5$), and weak predictors are those with an absolute value of the β coefficient less than 0.3 ($|\beta| < 0.3$).

As anticipated, a general linear model analysis demonstrated a strong positive association among all four spirometry parameters,

indicating a high degree of mutual correlation. The 6MWD emerged as a weak positive predictor for FEV1/FVC and FEF50%, but a strong positive predictor for FVC and FEV1. These findings are consistent with previous studies that have also reported a positive association between 6MWD and spirometry variables [23-25].

Moderately positive predictors of FEV1 and FEV1/FVC include DLCO%, PO₂, a-Sa(Hb)-O₂%, and t-Sa(Hb)-O₂%. Several studies have reported positive associations between DLCO% and spirometry parameters [26-29]. All these studies provide robust support for the positive relationships between DLCO% and various spirometry parameters, confirming the findings in this study. Moreover, lower DLCO% and lower FEV1 were associated with significantly increased morbidity in COPD patients compared to the reference group [30]. The positive relationship between oxygenation parameters [PO₂, a-Sa(Hb)-O₂%, and t-Sa(Hb)-O₂%] and spirometry values is a well-documented concept in the field of pulmonary medicine [23,31,32]. Additionally, FFMI was shown to be a weak positive predictor for all four spirometry outcomes, while




Figure 1. Standardized β coefficients with 95% confidence intervals for the effect of various predictors on forced vital capacity. Only statistically significant predictors are shown. CI, confidence interval; GOLD (I-IV), global initiative for chronic obstructive lung disease; BODE, body mass index/airflow obstruction/dyspnea/exercise capacity index; CAT, chronic obstructive pulmonary disease assessment test; mMRC, modified medical research council; SGQR-C, St. George's respiratory questionnaire; PO₂, arterial partial pressure of oxygen; DASS, depression anxiety stress scales; t-Sa(Hb)-O₂%, tissue hemoglobin oxygen saturation percentage; DLCO%, predicted diffusing FEV1, forced expiratory volume in one second; DLCO%, predicted diffusing capacity for carbon monoxide; FVC, forced vital capacity; a-Sa(Hb)-O₂%, arterial hemoglobin oxygen saturation percentage; FFMI, adjusted fat-free mass index; FEF50%, forced expiratory flow at 50% of FVC; 6MWD, 6-minute walk distance.



SMMI was a weak positive predictor for FEV1 and FEV1/FVC. There is considerable evidence that both FFMI and SMMI have a positive impact on spirometry outcomes [33-35], although their predictive strength can be weak, especially in the context of multifactorial influences in COPD patients.

BMI was identified as a weak positive predictor of FEV1 and FEV1/FVC. However, the β coefficients for BMI in relation to FVC and FEF50% were not statistically significant. These findings are consistent with our previously reported results, indicating that FEV1, FEV1/FVC, and DLCO values in pulmonary function tests were significantly higher in obese COPD patients compared to non-obese patients, whereas the FVC values were similar across all groups [21]. This outcome is commonly documented among individuals diagnosed with COPD [36-39].

Regarding negative predictors, strong negative predictors for spirometry outcomes include smoking-related factors, such as the number of years of smoking, pack-years, and smoking status. It is also evident that the BODE index and GOLD classification, relying extensively on FEV1 measurements, demonstrate a significant negative influence on spirometry parameters. Age and duration of illness have emerged as moderate negative predictors for spirometry readings, aligning with established findings in the field [40-42]. SGRQ-C, PCO₂, CAT, and mMRC were found to be weak predictors, consistent with previously published findings [43-46].

Factors such as the number of comorbidities, CCI, and PhA were found to have no significant impact on spirometry outcomes. Several studies have explored the relationship between these parameters and spirometry outcomes, yielding mixed results. For

Table 2. Predictors of forced vital capacity (FVC), forced expiratory volume in one second (FEV1), FEV1/FVC, and forced expiratory flow at 50% of FVC.

	FVC	FEV1	FEV1/FVC	FEF50%
Strong positive predictors	FEV1	FVC FEV1/FVC FEF50%	FEV1 FEF50%	FEV1 FVC FEV1/FVC
Moderate positive predictors	6MWD FEF50%	6MWD DLCO% PO ₂ t-Sa(Hb)-O ₂ % FFMI	DLCO% KCO% t-Sa(Hb)-O ₂ % PO ₂ a-Sa(Hb)-O ₂ %	—
Weak positive predictors	FFMI PO ₂ a-Sa(Hb)-O ₂ % FEV1/FVC DLCO% t-Sa(Hb)-O ₂ %	KCO% a-Sa(Hb)-O ₂ % SMMI BMI	FVC FFMI 6MWD BMI SMMI FVC	DLCO% PO ₂ KCO% t-Sa(Hb)-O ₂ % a-Sa(Hb)-O ₂ % 6MWD FFMI PhA
Strong negative predictors	Smoking years Pack-year Smoking status GOLD (I-IV)	GOLD (I-IV) BODE Pack-year Smoking status Smoking years	GOLD (I-IV) Smoking years Smoking status BODE Pack-year	Pack-year Smoking years Smoking status GOLD (I-IV)
Moderate negative predictors	BODE Age	mMRC Diagnosis SGRQ-C Age	Diagnosis mMRC SGRQ-C	BODE Diagnosis Age
Weak negative predictors	CAT mMRC SGRQ-C PCO ₂ DASS	CAT PCO ₂ Diagnosis DASS	CAT Age NRS-2002 PCO ₂ DASS	mMRC SGRQ-C CAT PCO ₂ NRS-2002
Not statistically significant	KCO% PhA CCI SMMI NRS-2002 BMI Comorbidities Diagnosis	NRS-2002 CCI Comorbidities PhA	CCI Comorbidities PhA	Comorbidities CCI SMMI DASS

FVC, forced vital capacity; FEV1, forced expiratory volume in one second; FEF50% and forced expiratory flow at 50% of FVC; 6MWD, 6-minute walk distance; DLCO%, predicted diffusing capacity for carbon monoxide; PO₂, arterial partial pressure of oxygen; t-Sa(Hb)-O₂%, tissue hemoglobin oxygen saturation percentage; FFMI, adjusted fat-free mass index; KCO%, predicted carbon monoxide transfer coefficient; a-Sa(Hb)-O₂%, arterial hemoglobin oxygen saturation percentage; SMMI, skeletal muscle mass index; BMI, body mass index; PhA, phase angle; GOLD (I-IV), global initiative for chronic obstructive lung disease; BODE, body mass index/airflow obstruction/dyspnea/exercise capacity index; mMRC, modified medical research council; SGRQ-C, St. George's respiratory questionnaire; CAT, chronic obstructive pulmonary disease assessment test; PCO₂, arterial partial pressure carbon dioxide; DASS, depression anxiety stress scales; CCI, Charlson comorbidity index; NRS-2002, nutritional risk screening.



instance, the number of comorbidities, as measured by tools such as the CCI, has been reported to have varying impacts on spirometry outcomes. While some research indicates no significant effect, others suggest correlations between higher comorbidity scores and poorer lung function [47,48]. PhA, a parameter derived from bioelectrical impedance analysis and often used as an indicator of nutritional status and overall health, has similarly been studied for its potential impact on lung function. Again, findings are mixed, with some studies indicating significant associations with spirometry variables, while others do not [49,50]. These discrepancies highlight the complexity of understanding how these factors influence respiratory health and suggest that further research is needed to clarify these relationships and their underlying mechanisms.

To further explore the issues being investigated, our objective was to develop a predictive model for the spirometry variables under consideration. Despite predictive models traditionally serving as indicators for the onset of illness, the development of predictive regression models among patients already affected by illness can prove highly advantageous. This approach holds significant potential for enhancing healthcare outcomes, optimizing resource management, and improving the effectiveness of medical interventions.

Particularly, our goal was to establish a predictive model using a reduced set of the most impactful predictors to select the best subset of predictors from a larger set of predictors. To achieve this, we employed stepwise regression, a method that sequentially selects predictors based on some statistical criteria. To mitigate bias in variable selection inherent to the stepwise method and to isolate the influence on each spirometry outcome from factors independent of the spirometry test itself, we initially excluded five predictors highly correlated with each spirometry variable. These predictors encompassed the remaining three spirometry outcomes, as well as the BODE and GOLD (I-IV) scores, which integrate FEV1 within their scoring systems.

The results of the linear regression model using the stepwise method are presented in Table 3. Indication I pertain to the stepwise regression conducted with standardized variables, aiming to determine comparable β coefficients. Positive predictors of FVC, FEV1, and FEV1/FVC are indicated in blue in the table, while negative predictors are highlighted in red. As expected, excluding five influential factors affecting the spirometry variables under investigation led to a decrease in the adjusted R-squared, which remains around 0.65 across the three models presented. The R-squared for modelling FEF50% was below 0.5, thus rendering this analysis unreliable, and it has been excluded. In Table 3, indication II denotes the results of stepwise regression analysis conducted on unstandardized variables, aimed at establishing practical regression equations (1), (2), and (3), which can be found in the *Supplementary Material*.

After excluding five spirometry-related parameters, FFMI and 6MWD emerge as the stronger positive predictors for all three spirometry readings considered in the modelling. Other positive predictors include DLCO% and a-Sa(Hb)-O₂%. PO₂ is a statistically significant positive predictor for FVC, while BMI is a significant positive predictor for FEV1 and FEV1/FVC. The strongest negative predictor of spirometry variables is smoking years, with other statistically significant negative predictors including age, illness duration, mMRC, CAT, PCO₂, and SGQR-C.

To mitigate the complexity and intricacy associated with data processing and foster a more profound comprehension of interre-

lationships among variables within a dataset, factorial analysis is undertaken. Based on the scree plot (see *Supplementary Material*), it has been demonstrated that selecting five factors represents the optimal choice. This selection ensures a balanced representation of underlying structures while effectively capturing a satisfactory amount of variance (62%) within the dataset. After the initial extraction of factors in factor analysis, varimax rotation is employed to simplify and enhance the interpretability of the factor structure by maximizing the variance of squared loadings on each factor and ensuring orthogonality among factors.

Rotated factor loadings and communalities following varimax rotation are displayed in Table 4. The highest factor loadings for Factor 1 include spirometry parameters, BODE, GOLD (I-IV), a-Sa(Hb)-O₂%, t-Sa(Hb)-O₂%, PO₂, PCO₂, DLCO%, and KCO%. Based on this, Factor 1 can be concluded to be associated with lung function. Factor 2 exhibits the highest loadings for CAT, DASS, mMRC, SGRQ-C, and NRS-2002, broadly reflecting the health status of the patients. Factor 3 is characterized by the highest loadings for FFMI, SSMI, BMI, and PhA, indicating its association with nutritional status. Factor 4 appears to be linked to age, as reflected by the highest loadings of age, illness duration, CCI, and 6MWD. Lastly, Factor 5 clearly pertains to smoking, with the highest loadings for pack-years, smoking years, and smoking status. Ultimately, factor analysis divided the 29 variables under consideration into five factors, respectively related to lung function (Factor 1), health status (Factor 2), nutritional status (Factor 3), age (Factor 4), and smoking (Factor 5). The highest loadings for each factor are highlighted in bold in Table 4, with different colours indicating each factor (Factor 1 - black, Factor 2 - blue, Factor 3 - red, Factor 4 - green, Factor 5 - purple).

Following factorial analysis, four new variables were generated: health status (Factor 2), nutritional status (Factor 3), age (Factor 4), and smoking (Factor 5). This was achieved by combining individual variables with the highest loadings within each factor. Specifically, each given variable was multiplied by its respective loading; the products of the loadings and variable values were summed and then divided by the sum of the loadings, considering the sign of the loadings. This procedure was undertaken to create a new model for three pulmonary variables (FVC, FEV1, FEV1/FVC), with the four newly created variables serving as predictors. The variables from Factor 1 were not consolidated into a new variable since they exhibit strong intercorrelations and are related to lung function, like the variables for which the model is being developed.

The results of the multiple linear regression with the four newly formed predictors are presented in Table 3, indicated by indicator III. As can be observed, for all three spirometry variables, the results are consistent — nutritional status (Factor 3) is a positive predictor of spirometry outcomes, while poorer health status, age, and smoking are negative predictors. It is important to note that in mathematical terms, an increase in the health status variable (mMRC, CAT, DASS, SGRQ-C, and NRS-2002) corresponds to a decrease in the dependent variables (FVC, FEV1, FEV1/FVC).

In conclusion, following the factorial analysis and multiple linear regression conducted with the four newly formed factors, it can be stated that spirometry outcomes are most significantly influenced by smoking (negatively), followed by poor health status (negatively), nutritional status (positively), and finally age (negatively). The regression equations for FVC (4), FEV1 (5), and FEV1/FVC (6) can be found in the *Supplementary Material*.



Table 3. The results of statistical modeling using stepwise regression and multiple linear regression following factorial analysis.

Stat. Method*	Indep. Var.	Dep. Var.**	β	SE	p	R ² adj (%)	Regr. Equat.				
I	FVC	FFMI	0.358	0.060	0.000	67.50					
		6MWD	0.280	0.037	0.000						
		PO ₂	0.212	0.051	0.000						
		DLCO%	0.145	0.052	0.005						
		a-Sa(Hb)-O ₂ %	0.079	0.034	0.020						
		mMRC	-0.116	0.035	0.001						
		CAT	-0.122	0.036	0.001						
		Age	-0.147	0.036	0.000						
		SMMI	-0.167	0.035	0.000						
		PCO ₂	-0.168	0.033	0.000						
		Smoking years	-0.558	0.034	0.000						
							(1)				
I	FEV1	FFMI	0.243	0.046	0.000	67.76					
		6MWD	0.216	0.054	0.000						
		DLCO%	0.175	0.039	0.000						
		a-Sa(Hb)-O ₂ %	0.134	0.038	0.000						
		BMI	0.104	0.053	0.014						
		Diagnosis	-0.093	0.039	0.018						
		mMRC	-0.095	0.046	0.041						
		CAT	-0.113	0.046	0.014						
		Age	-0.114	0.037	0.002						
		PCO ₂	-0.134	0.034	0.000						
		SGRQ-C	-0.146	0.039	0.000						
		Smoking years	-0.387	0.036	0.000						
									(2)		
I	FEV1/FVC	FFMI	0.233	0.047	0.000	65.37					
		t-Sa(Hb)-O ₂ %	0.199	0.046	0.000						
		6MWD	0.179	0.053	0.000						
		BMI	0.168	0.043	0.000						
		DLCO%	0.122	0.076	0.027						
		a-Sa(Hb)-O ₂ %	0.112	0.044	0.010						
		Diagnosis	-0.085	0.039	0.032						
		SGRQ-C	-0.132	0.053	0.012						
		DASS	-0.153	0.046	0.001						
		Age	-0.239	0.057	0.000						
		Smoking years	-0.288	0.042	0.000						
									(3)		
		III	FVC	Factor 3	0.132			0.015	0.014	60.81	
Factor 4	-0.138			0.012	0.002						
Factor 2	-0.145			0.011	0.000						
Factor 5	-0.242			0.018	0.000						
							(4)				
III	FEV1	Factor 3	0.157	0.016	0.000	59.24					
		Factor 4	-0.126	0.012	0.037						
		Factor 2	-0.174	0.011	0.000						
		Factor 5	-0.220	0.019	0.000						
							(5)				
III	FEV1/FVC	Factor 3	0.170	0.017	0.000	54.84					
		Factor 4	-0.123	0.020	0.042						
		Factor 2	-0.182	0.012	0.000						
		Factor 5	-0.207	0.013	0.000						
							(6)				

Factor 2, HEALTH STATUS; Factor 3, NUTRITION STATUS; Factor 4, AGE; Factor 5, SMOKING. *Parameters strongly related to the independent variable [the rest of the three spirometry parameters, BODE, and GOLD (I-IV)] have been excluded in regression models as predictors; **only statistically significant predictors are shown. I, stepwise regression with standardized variables; II, stepwise regression with unstandardized variables; III, multiple linear regression (using four new parameters from factorial analysis as predictors) with standardized variables; IV, multiple linear regression (using four new parameters from factorial analysis as predictors) with unstandardized variables. STAT. METHOD, statistical method; INDEP. VAR., independent variable; DEP. VAR., dependent variable; SIGNIF. PREDIC., predictor with statistically significant β; p, p-value; SE, standard error for β; REGR. EQUAT., multiple regression equation; FVC, forced vital capacity; FFMI, adjusted fat-free mass index; 6MWD, 6-minute walk distance; PO₂, arterial partial pressure of oxygen; DLCO%, predicted diffusing capacity for carbon monoxide; a-Sa(Hb)-O₂%, arterial hemoglobin oxygen saturation percentage; mMRC, modified medical research council; CAT, chronic obstructive pulmonary disease assessment test; SMMI, skeletal muscle mass index; PCO₂, arterial partial pressure carbon dioxide; FEV1, forced expiratory volume in one second; SGRQ-C, St. George's respiratory questionnaire.



Table 4. Rotated factor loadings and communalities (varimax rotation).

	Variable	Factor 1	Factor 2	Factor 3	Factor 4	Factor 5	Communality
1	Age	-0.026	-0.133	-0.017	0.852	0.042	0.747
2	BMI	-0.007	-0.038	0.844	0.012	-0.140	0.733
3	Smoking status	0.162	0.034	-0.129	-0.411	0.533	0.497
4	Smoking years	-0.023	0.033	-0.009	0.094	0.782	0.743
5	Pack-year	-0.096	0.044	0.078	0.122	0.856	0.644
6	Diagnosis	-0.109	0.194	-0.100	0.384	-0.131	0.224
7	mMRC	-0.390	0.694	-0.007	0.108	0.076	0.652
8	CAT	-0.215	0.757	-0.008	0.030	-0.038	0.621
9	GOLD (I-IV)	-0.776	0.206	-0.087	0.087	0.045	0.661
10	FVC	0.520	-0.115	0.084	-0.261	0.390	0.450
11	FEV1	0.834	-0.172	0.093	-0.090	-0.050	0.745
12	FEV1/FVC%	0.714	-0.114	0.200	-0.106	-0.219	0.621
13	FEF50%	0.539	-0.116	0.101	-0.172	-0.157	0.368
14	DLCO%	0.496	-0.218	0.445	0.017	-0.287	0.575
15	KCO%	0.345	-0.203	0.482	0.207	-0.344	0.554
16	t-Sa(Hb)-O ₂ %	0.577	-0.050	-0.054	-0.095	0.030	0.348
17	a-Sa(Hb)-O ₂ %	0.705	0.117	-0.073	0.188	0.121	0.566
18	PO ₂	0.699	0.118	-0.048	0.122	0.120	0.534
19	PCO ₂	-0.339	0.066	0.131	-0.016	-0.113	0.149
20	6MWD	0.371	-0.389	-0.068	-0.407	0.017	0.459
21	Comorbidities	-0.013	0.265	-0.518	-0.246	0.040	0.401
22	DASS	0.045	0.758	-0.055	0.017	0.056	0.583
23	SGRQ-C	-0.338	0.690	0.006	0.027	0.062	0.596
24	NRS-2002	0.128	0.634	0.041	-0.107	-0.018	0.431
25	BODE	-0.659	0.469	-0.182	0.201	0.054	0.730
26	CCI	0.006	-0.099	0.043	0.776	0.046	0.617
27	SMMI	-0.047	0.002	0.868	-0.039	0.064	0.761
28	PhA	0.054	0.065	0.353	-0.115	0.078	0.151
29	FFMI	-0.048	-0.015	0.869	0.004	0.039	0.759
	Variance	5.058	3.290	3.002	2.298	2.207	15.855
	% Variance	0.184	0.133	0.124	0.091	0.088	0.616

Factor 1, lung function; factor 2, health status (questionnaires and tests - higher values indicate poorer health status); factor 3, nutritional status; factor 4, age; factor 5, smoking. BMI, body mass index; mMRC, modified medical research council; CAT, chronic obstructive pulmonary disease assessment test; GOLD (I-IV), global initiative for chronic obstructive lung disease; FVC, forced vital capacity; FEV1, forced expiratory volume in one second; FEF50%, forced expiratory flow at 50% of FVC; DLCO%, predicted diffusing capacity for carbon monoxide; KCO%, predicted carbon monoxide transfer coefficient; t-Sa(Hb)-O₂%, tissue hemoglobin oxygen saturation percentage; a-Sa(Hb)-O₂%, arterial hemoglobin oxygen saturation percentage; PO₂, arterial partial pressure of oxygen; PCO₂, arterial partial pressure carbon dioxide; 6MWD, 6-minute walk distance; DASS, depression anxiety stress scales; SGRQ-C, St. George's respiratory questionnaire; NRS-2002, nutritional risk screening; BODE, body mass index/airflow obstruction/dyspnea/exercise capacity index; CCI, Charlson comorbidity index; SMMI, skeletal muscle mass index; PhA, phase angle; FFMI, adjusted fat-free mass index.

Conclusions

In the current study, encompassing a large cohort of 534 COPD patients from five distinct centers in Croatia, the prediction of spirometry functional test parameters (specifically FVC, FEV1, FEV1/FVC, and FEF50%) was conducted, involving 29 different variables. This prediction was achieved through the application of several statistical methods, including a general linear model, step-wise regression, and subsequent multiple regression models, followed by the results of factorial analysis.

Although the findings of the study align with established knowledge regarding factors influencing spirometry pulmonary test parameters in patients diagnosed with COPD, it is important to note that this study precisely determined the magnitude of these vari-

ables' impacts, as well as predictive models for spirometry outcomes in patients already diagnosed with COPD.

The study also demonstrated that smoking has the most detrimental effect on spirometry variables, followed by poor health status, and then the age of the patients. Good nutritional status positively affects spirometry variables. However, patients with a higher BMI exhibit better FEV1 and FEV1/FVC outcomes as has been documented in a considerable number of studies.

References

1. Global Initiative for Chronic Obstructive Lung Disease. Global strategy for the diagnosis, management and prevention of COPD. 2021 report. Available from: <https://goldcopd.org>.



- org/wp-content/uploads/2020/11/GOLD-REPORT-2021-v1.1-25Nov20_WMV.pdf.
- Halpin DMG, Celli BR, Criner GJ, et al. The GOLD Summit on chronic obstructive pulmonary disease in low- and middle-income countries. *Int J Tuberc Lung Dis* 2019;23:1131-41.
 - Lopez-Campos JL, Tan W, Soriano JB. Global burden of COPD. *Respirology* 2016;21:14-23.
 - GBD 2017 Causes of Death Collaborators. Global, regional, and national age-sex-specific mortality for 282 causes of death in 195 countries and territories, 1980-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet* 2018;392:1736-88.
 - Pellegrino R, Viegi G, Brusasco V, et al. Interpretative strategies for lung function tests. *Eur Respir J* 2005;26:948-68.
 - Miller MR, Hankinson J, Brusasco V, et al. Standardisation of spirometry. *Eur Respir J* 2005;26:319-38.
 - Global Initiative for Chronic Obstructive Lung Disease. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease. 2023 report. Available from: <https://goldcopd.org/2023-gold-report-2/>.
 - Miller MR, Crapo R, Hankinson J, et al. General considerations for lung function testing. *Eur Respir J* 2005;26:153-61.
 - Guerra S, Sherrill DL, Kurzius-Spencer, M, et al. The course of persistent airflow limitation in subjects with and without asthma. *Respir Med* 2008;102(10):1473-82.
 - Hankinson JL, Odencrantz JR, Fedan KB. Spirometric reference values from a sample of the general U.S. population. *Am J Respir Crit Care Med* 1999;159:179-87.
 - Holguin F, Folch E, Redd SC, Mannino DM. Comorbidity and mortality in COPD-related hospitalizations in the United States, 1979 to 2001. *Chest* 2017;128:2005-11.
 - Pellegrino R, Viegi G, Brusasco V. Interpretative strategies for lung function tests. *Eur Respir J* 2005;26:948-68.
 - Stanojevic S, Wade A, Stocks J, et al. Reference ranges for spirometry across all ages: a new approach. *Am J Respir Crit Care Med* 2008;177:253-60.
 - Miura S, Iwamoto H, Omori K, et al. Accelerated decline in lung function in adults with a history of remitted childhood asthma. *Eur Respir J* 2022;59(1):2100305-15.
 - Quanjer PH, Stanojevic S, Cole TJ, et al. Multi-ethnic reference values for spirometry for the 3-95-yr age range: the global lung function 2012 equations. *Eur Respir J* 2012;40:1324-43.
 - Swanney MP, Jensen RL, Crichton DA, et al. FEV(6) is an acceptable surrogate for FVC in the spirometric diagnosis of airway obstruction and restriction. *Am J Respir Crit Care Med* 2000;162:917-9.
 - Burney PG, Laitinen LA, Perdrizet S, et al. Validity and repeatability of the IUATLD (1984) bronchial symptoms questionnaire: an international comparison. *Eur Respir J* 1989;2:940-5.
 - Sin DD, McAlister FA, Man SF, Anthonisen NR. Contemporary management of chronic obstructive pulmonary disease: scientific review. *JAMA* 2003;290:2301-12.
 - Vogelmeier CF, Criner GJ, Martinez FJ, et al. Global strategy for the diagnosis, management, and prevention of chronic obstructive lung disease 2017 report. GOLD executive summary. *Am J Respir Crit Care Med* 2017;195:557-82.
 - Muršić D, Jalušić Glunčić T, Ostojić J, et al. Differences in nutritional status of patients with chronic obstructive pulmonary disease between Mediterranean and non-Mediterranean regions in Croatia. *Monaldi Arch Chest Dis* 2023;93:2667. DOI: 10.4081/monaldi.2023.2667.
 - Muršić D, Jalušić Glunčić T, Ostojić J, et al. Body composition, pulmonary function tests, exercise capacity, and quality of life in chronic obstructive pulmonary disease patients with obesity. *Postgrad Med J* 2024;100:469-74.
 - Kyle UG, Bosaeus I, De Lorenzo AD, et al. Bioelectrical impedance analysis—part I: review of principles and methods. *Clin Nutr* 2004;23:1226-43.
 - Pinto-Plata VM, Cote C, Cabral H, et al. The 6-min walk distance: change over time and value as a predictor of survival in severe COPD. *Eur Respir J* 2004;23:28-33.
 - Spruit MA, Polkey MI, Celli B, et al. Predicting outcomes from 6-minute walk distance in chronic obstructive pulmonary disease. *J Am Med Dir Assoc* 2012;13:291-7.
 - Pitta F, Troosters T, Spruit MA, et al. Characteristics of physical activities in daily life in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2005;171:972-7.
 - Graham BL, Brusasco V, Burgos F, et al. 2017 ERS/ATS standards for single-breath carbon monoxide uptake in the lung. *Eur Respir J* 2017;49:1600016.
 - Li Y, Li XY, Yuan LR, et al. Evaluation of small airway function and its application in patients with chronic obstructive pulmonary disease (Review). *Exp Ther Med* 2021;22:1386-94.
 - Abd-Allah HM, Abdalla H, Ahmed S, Mohammed MH. Role of single breath diffusing capacity for carbon monoxide in patients with chronic obstructive pulmonary disease. *Al-Azhar Assiut Med J* 2021;19:61-7.
 - Choi J, Sim JK, Oh JY, et al. Prognostic marker for severe acute exacerbation of chronic obstructive pulmonary disease: analysis of diffusing capacity of the lung for carbon monoxide (DLCO) and forced expiratory volume in one second (FEV1). *BMC Pulm Med* 2021;21:152.
 - Celli BR, Cote CG, Marin JM, et al. The body-mass index, airflow obstruction, dyspnea, and exercise capacity index in chronic obstructive pulmonary disease. *N Engl J Med* 2004;350:1005-12.
 - Viegi G, Pedreschi M, Pistelli F, et al. Prevalence of airways obstruction in a general population: European Respiratory Society vs American Thoracic Society definition. *Chest* 2000;117:330-45.
 - Schols AM, Broekhuizen R, Weling-Scheepers CA, Wouters EF. Body composition and mortality in chronic obstructive pulmonary disease. *Am J Clin Nutr* 2005;82:53-9.
 - Marquis K, Debigaré R, Lacasse Y, et al. Midthigh muscle cross-sectional area is a better predictor of mortality than body mass index in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2002;166:809-13.
 - Vestbo J, Prescott E, Almdal T. Body mass, fat-free body mass, and prognosis in patients with chronic obstructive pulmonary disease from a random population sample: findings from the Copenhagen City Heart Study. *Am J Respir Crit Care Med* 2006;173:79-83.
 - Maltais F, Decramer M, Casaburi R, et al. An official American Thoracic Society/European Respiratory Society statement: update on limb muscle dysfunction in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2014;189:e15-62.
 - Lando C, Prescott E, Lange P, et al. Prognostic value of nutritional status in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1999;160:1856-61.
 - Spelta F, Fratta Pasini AM, Cazzoletti L, et al. Body weight and mortality in COPD: focus on the obesity paradox. *Eat Weight Disord* 2018;23:15-22.



38. Schols AM, Slangen J, Volovics L, Wouters EF. Weight loss is a reversible factor in the prognosis of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1998;157:1791-7.
39. Casanova C, de Torres JP, Aguirre-Jaime A, et al. The body mass index, airflow obstruction, dyspnea, and exercise capacity index in chronic obstructive pulmonary disease. *Chest*. 2005;128:911-9. DOI: 10.1378/chest.128.2.911
40. Knudson RJ, Lebowitz MD, Holberg CJ, Burrows B. Changes in the normal maximal expiratory flow-volume curve with growth and aging. *Am Rev Respir Dis* 1983;127:725-34.
41. Lange P, Celli B, Agustí A, et al. Lung-function trajectories leading to chronic obstructive pulmonary disease. *N Engl J Med* 2015;373:111-22.
42. Jones PW, Harding G, Berry P, et al. Development and first validation of the COPD Assessment Test. *Eur Respir J* 2009;34:648-54.
43. Pauwels RA, Buist AS, Calverley PM, et al. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: NHLBI/WHO global initiative for chronic obstructive lung disease (GOLD) workshop summary. *Am J Respir Crit Care Med* 2001;163:1256-76.
44. Cazzola M, Macklem PT, Stolz D. Unmet needs in COPD: a vision for the future. *Int J Chron Obstruct Pulmon Dis* 2008; 3:411-29.
45. de Torres JP, Casanova C, Hernandez C, et al. Gender and COPD in patients attending a pulmonary clinic. *Chest* 2005;128:2012-6.
46. Wan ES, Balte P, Schwartz JE, et al. Association between preserved ratio impaired spirometry and clinical outcomes in US adults. *JAMA* 2021;326:2287-98.
47. Zhang N, Lin Q, Jiang H, et al. Age-adjusted Charlson Comorbidity Index as effective predictor for in-hospital mortality of patients with cardiac arrest: a retrospective study. *BMC Emerg Med* 2023;23:7.
48. Quan H, Li B, Couris CM, et al. Updating and validating the Charlson Comorbidity Index and score for risk adjustment in hospital discharge abstracts using data from 6 countries. *Am J Epidemiol* 2011;173:676-82.
49. Valisoltani N, Mohammadi H, Aliannejad R, et al. Association of phase angle with sarcopenia and muscle function in patients with COPD: a case-control study. *BMC Pulm Med* 2024;24:18.
50. Ruiz-Margáin A, Xie JJ, Román-Calleja BM, et al. Phase angle from bioelectrical impedance for the assessment of sarcopenia in cirrhosis with or without ascites. *Clin Gastroenterol Hepatol* 2021;19:1941-9.e2.

Online supplementary material:

Table S1. Predictors of forced vital capacity.

Table S2. Predictors of forced expiratory volume in one second.

Table S3. Predictors of forced expiratory volume in one second/forced vital capacity ratio.

Table S4. Predictors of forced expiratory flow at 50% of forced vital capacity.

Figure S1. Patient flow chart.

Figure S2. Standardized β coefficients with 95% confidence intervals for the effect of various predictors on FEV1. Only statistically significant predictors are shown.

Figure S3. Standardized β coefficients with 95% confidence intervals for the effect of various predictors on FEV1/FVC. Only statistically significant predictors are shown.

Figure S4. Forced expiratory volume in one second. Standardized β coefficients with 95% confidence intervals for the effect of various predictors on FEF50%. Only statistically significant predictors are shown.

Supplementary material. Computer outcomes for forced vital capacity in relation to predictor, age and sex.

Received: 25 June 2024; Accepted: 20 September 2024; Early view: 15 November 2024.

Contributions: Erim Bešić, Davorka Muršić, Miroslav Samaržija, Andrea Vukić Dugac: conception and design of the article and critical revision for important intellectual content. Davorka Muršić, Tajana Jalušić Glunčić, Jelena Ostojčić, Sanda Škrinjarić-Cincar, Martina Dokoza, Nataša Karamarković Lazarušić, Andrea Vukić Dugac: acquisition of the data and critical revision of the article for important intellectual content. Erim Bešić, Davorka Muršić, Andrea Vukić Dugac: acquisition, analysis, and interpretation of the data, and drafting of the article. All authors critically revised the manuscript, agreed to be fully accountable for ensuring the integrity and accuracy of the work, and read and approved the final manuscript.

Conflict of interest: the authors declare that they have no competing interests, and all authors confirm accuracy.

Ethics approval and consent to participate: the study protocol was approved by the Ethical Review Committee of the University Hospital Centre Zagreb (ID 02/18 AG) as well as by the Ethical Committee of the University Hospital Centre Rijeka, the University Hospital Centre Osijek, and Zadar General Hospital.

Informed consent: written informed consent was obtained from a legally authorized representative(s) for anonymized patient information to be published in this article. The manuscript does not contain any individual person's data in any form.

Patient consent for publication: written informed consent was obtained from a legally authorized representative(s) for anonymized patient information to be published in this article. The manuscript does not contain any individual person's data in any form.

Availability of data and materials: data and materials are available from the corresponding author upon request.

Publisher's note: all claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).

