

Dilated cardiomyopathy - guidelines for personalization of care

Antonella Spinelli, Gaetano Marino, Alfredo Macchiusi, Stefania Angela Di Fusco, Carlo Pignalberi, Furio Colivicchi

Clinical and Rehabilitation Cardiology Unit, San Filippo Neri Hospital - ASL Roma 1, Rome, Italy

Abstract

The term “dilated cardiomyopathy” (DCM) refers to a family of diseases characterized by complex interactions between environment and genetic predisposition. Diagnostic tools such as cardiac magnetic resonance imaging should be systematically implemented in clinical practice to define the etiological cause and undertake a specific treatment. We present the case of a young man with DCM and severe left ventricular dysfunction. The patient has a family history of sudden cardiac death (SCD). He is affected by a psychiatric pathology and has a history of alcohol and drug addiction. Many diagnostic hypotheses have been considered for etiological research. The use of a wearable defibrillator to temporarily protect the patient from SCD risk pending the completion of the diagnostic path to establish DCM etiology and the implementation of optimal medical therapy was considered. However, because of the underlying psychiatric pathology and the possibility of poor adherence to the maintenance of the wearable device, the patients would not have been sufficiently protected. Due to the young age of the patient, it was decided to make a personalized therapeutic choice, and he underwent implantation of a subcutaneous defibrillator.

Key words: heart failure, subcutaneous defibrillator.

Correspondence to: Antonella Spinelli, Clinical and Rehabilitation Cardiology Unit, San Filippo Neri Hospital ASL Roma 1, via Giovanni Martinotti 20, 00135, Rome, Italy.

Tel: 339-6257587. E-mail: antonellaspinelli81@gmail.com

Introduction

Cardiomyopathy is defined as a “myocardial disorder in which the heart muscle is structurally and functionally abnormal, in the absence of coronary artery disease (CAD), hypertension, valvular disease, and congenital heart disease (CHD) sufficient to cause the observed myocardial abnormality” [1]. Patients with reduced ejection fraction (EF) have a high risk of sudden cardiac death (SCD), so it is imperative to protect them throughout their diagnostic examination and treatment course aimed at left ventricular function recovery. The early implantation of an implantable defibrillator (ICD) to prevent SCD is often debated in the clinical management of these patients [1].

Case Report

We report the case of a 28-year-old patient with a high school education and a family history of sudden death (a maternal uncle deceased at age 39 and a sister with Rett syndrome died in her sleep at age 18). The patient practiced sports at a competitive level until the age of 12. He had a clinical history of alcohol abuse since age 14, regular intake of cannabinoids and occasional intake of cocaine, and a previous diagnosis of attention-deficit/hyperactivity disorder (ADHD). The patient was treated in a psychiatric diagnostic and treatment service with the following therapy: gabapentin, quetiapine, clonazepam, atomoxetine hydrochloride, oxcarbazepine, and

trazodone hydrochloride. In September 2022, during his stay at an addiction cessation facility, he had the onset of a dry cough and worsening dyspnea and two subsequent febrile episodes in November 2022. In February 2023 he accessed the emergency room for cardio-respiratory failure after an accidental fall.

On admission, the patients had severe left ventricular global systolic dysfunction with a left ventricular EF of 10% in the absence of regional wall-motion abnormalities. The left ventricular telediastolic volume (LVEDV) was 251 mL/m², and the left ventricular telesystolic volume (LVESV) was 197 mL/m². He had no significant valvulopathy and the right ventricle was enlarged but normokinetic. Blood pressure was 100/60 mmHg and oxygen saturation was 91%. The electrocardiogram showed a sinus rhythm at 110 bpm with mild nonspecific intraventricular conduction disturbance (QRS 112 msec) and nonspecific ventricular repolarization abnormalities. Procalcitonin was negative (0.24 ng/mL, n.v. 0.0 - 0.5), hemoglobin (Hb) levels, white blood cells (WBC), and platelet (PLT) count were in a normal range (Hb 12.0 g/dL, n.v. 12.0-16.0; WBC 9.100/uL, n.v. 4.0-11.0; PLT 139000/uL n.v. 130000-400000); peak C-reactive protein (CRP) was 7.93 mg/dL; peak ultrasensitive troponin was 52.50 ng/L; and peak N-terminal pro B-type natriuretic peptide was 1297 pg/mL. Blood tests were done to study alcohol-related damage to the heart.

Tests for liver function were normal. Tests for heavy alcohol use (anemia, thrombocytopenia, folate, vitamin B12 deficiency, electrolyte alterations, and desialyted transferrin) were normal too. The chest radiography confirmed the clinical suspicion of bilateral pneu-



monia complicated by bilateral pleural effusion. Pneumonia could be a further cause of decompensation. Antibiotic therapy was started with subsequent normalization of CRP. The research of cardiopneumotropic virus antibodies showed immunoglobulin G (IgG) positivity for cytomegalovirus (71 U/mL) and parvovirus B19 (4.90 U/mL), whereas specific immunoglobulin M antibodies were negative. A coronarography excluded significant coronary stenosis. Cardiac magnetic resonance imaging (MRI) performed 11 days after admission confirmed a severe systolic dysfunction (EF 14%, LVEDV: 320 mL/m², LVESV: 275 mL/m²) with a reduced thickness of the midventricular antero-septal wall. Furthermore, the left ventricular myocardial wall had areas of altered signal, hyperintense on T2W sequences, suggestive of myocardial edema, associated with signal hyperintensity of the left periventricular adipose tissue, which showed marked late enhancement immediately after intravenous contrast administration (Figure 1).

Symptomatic sustained monomorphic ventricular tachycardia lasting 40 seconds was observed during hospitalization in the intensive care unit (ICU). It was self-limited and no further episodes of this type were recorded. During the remaining days of hospitalization, the finding of thrombocytosis, probably because of infectious or inflammatory processes, required therapy with acetylsalicylic acid. We have observed a progressive clinical improvement, although in the absence of recovery of EF, it is indicative of a persistent significant risk of life-threatening arrhythmias. Therefore, pending the completion of diagnostic examinations (genetic testing results) and pharmacological therapy titration before discharge, the use of wearable defibrillators (WCD) was considered. However, the psychiatric consultant pointed out the risk of poor adherence to the proper use of the device given the risk of impulsive behaviors related to the underlying psychiatric disorder, ADHD, and the history of alcohol and drug




Figure 1. Images showing diffuse enhancement of adipose tissue and subepicardial myocardium throughout the left ventricle wall (A-C). The hyperintense signal on T2W of the left ventricular myocardial wall is suggestive of edema, associated with hyperintense signal of the left periventricular adipose tissue (D).

abuse. After a comprehensive interview with the patient and his family, a multidisciplinary team, including a clinical cardiologist, electrophysiologist, and psychiatrist, agreed with the indication of subcutaneous defibrillator implantation (S-ICD).

The implantation procedure was performed in the absence of complications. The patient was discharged after 18 days of hospital stay, in satisfactory clinical-hemodynamic compensation, with the following pharmacological prescriptions: dapagliflozin 10 mg/day, sacubitril/valsartan 24/26 mg×2/day, furosemide 25 mg/day, canrenone 100 mg/day, bisoprolol 5 mg×2/day, and acetylsalicylic acid 100 mg/day, and the psychiatric therapy previously used was confirmed. At outpatient follow-up after 1 month from discharge, the EF was unchanged (15%). Acetylsalicylic acid was suspended for normalization of PLT count. Pharmacological therapy was not up-titrated due to the finding of low arterial pressure values. At 3 months, EF 28% was shown and cardiac MRI, performed within the expected time frame after ICD implantation (after at least 6-8 weeks from presentation), was affected by the presence of gross artifacts due to the presence of ferromagnetic material, so it was considered inconclusive (Figure 2).

Genetic evaluation results did not identify variants with probable or possible pathogenic significance for major dilated cardiomyopathy (DCM)-associated genes. Therefore, a pathway for the possible need for cardiac transplantation/LVAD implantation was also considered. Indeed, the patient reported complete abstinence from drugs since discharge, confirmed by negativity on toxicological testing. However, important concerns remained regarding psychiatric issues related to ADHD and the relatively short period of drug abstinence. It was decided to continue monthly follow-ups to intercept any changes in both the cardiological and psychiatric picture. At the follow-up performed in October 2023, EF improved and left ventricular volumes were reduced (EF 40%, LVEDV 98 mL/m², and LVESV 59 mL/m²).




Figure 2. Cardiac magnetic resonance imaging performed 8 weeks after subcutaneous defibrillator implantation.

Discussion

We report the case of a patient with non-ischemic DCM and severe left ventricular contractile dysfunction. Previous studies have confirmed that there are many known and unknown causes of DCM; the known causes are mainly infection, genetic, and immune factors. Studies have confirmed that viral myocarditis (VMC) is one of the possible causes for DCM. The molecular mechanisms underlying the progression from VMC to DCM remain unclear and require further investigation. It is likely that acute inflammation may progress to subacute and chronic stages and finally to tissue remodeling, fibrosis, and loss of myocardium architecture and contractile function. The latter chronic damages lead to the development of DCM [1,2]. According to literature data, the estimated prevalence of DCM is about 1:2500 individuals in the general population, slightly higher in men, with a female-to-male ratio between 1:1.3 and 1:1.5 [3,4]. In our patient, the presumptive etiological hypothesis is that DCM may be secondary to VMC, which probably occurred in November 2022 during febrile episodes. Overall, between 0 and 52% of patients with acute myocarditis develop DCM [4,5].

Retrospective studies report that 9-16% of unexplained non-ischemic DCM cases have histological evidence of myocarditis. Other studies estimate a wider range of 10-50% of DCM cases having evidence of myocarditis as could be determined by endomyocardial biopsy [5,6]. Cardiac MRI findings, showing features typical of myocardial fibrosis, supported the hypothesis and provided valuable diagnostic and arrhythmic risk stratification information, in agreement with recent European Society of Cardiology (ESC) guidelines (recommendation class I - level of evidence B). Indeed, the presence of late gadolinium enhancement, observed in 25-30% of patients with DCM, is a strong predictor of ventricular arrhythmias and all-cause mortality [1,7]. We performed genetic evaluation, which is recommended as Class I by the 2022 ESC guidelines on heart failure management and was confirmed as Class I in the most recent 2023 ESC guidelines on cardiomyopathies [1,7,8].

A panel of genes associated with DCM was subjected to analysis: *MYBPC3*, *MYH7*, *TNNT2*, *TNNI3*, *TPM1*, *ACTC1*, *MYL2*, *MYL3*, *CSRP3*, *TNNC1*, *PNL*, *LAMP2*, *GLA*, *TTR*, *PRKAG2*, *TNNT2*, *TNNI3*, *SCN5A* and *LMNA*. Genetic evaluation results did not identify variants with probable or possible pathogenic significance for major DCM-associated genes. The genetic test for Rett syndrome was not considered useful by the geneticist for several reasons: it was negative in both screened parents; Rett syndrome is almost exclusive to females and in males it is lethal in the first few years of life or during fetal life. The definitive diagnosis of VMC requires an endomyocardial biopsy, but in clinical practice biopsy is rarely used. In our case, the anamnestic reference (recent flu), the finding of IgG for CMV and parvovirus B19 made the hypothesis of viral origin very probable. Cardiac MRI also showed myocardial edema and fibrosis. According to the most recent guidelines, endomyocardial biopsy is recommended when other methods are insufficient [9]. Other etiological hypotheses that we have considered, included DCM from alcohol abuse, accounting for up to 32% of cases of DCM, or from toxic substance abuse (cocaine and/or antipsychotic drugs) [3,10,11]. Alcohol-induced cardiomyopathy (AC) is a type of heart disease caused by long-term alcohol use. The amount of alcohol needed to cause AC is usually thought to be more than 80 grams a day over 5 years, but this is still debated. AC develops in different ways. Alcohol and a metabolite of alcohol called acetaldehyde can cause it. Genes also play a part. Not all heavy drinkers have DCM. AC does not have any specific cardiac signs or

Table 1. Recommendations for an implantable or wearable cardioverter defibrillator in primary prevention for patients with dilated cardiomyopathy in agreement with 2023 European Society of Cardiology (ESC) guidelines for the management of cardiomyopathies [1], and 2021 ESC guidelines for the diagnosis and treatment of acute and chronic heart failure [16].

Recommendations	Class	Level
An ICD should be considered to reduce the risk of sudden death and all-cause mortality in patients with DCM, symptomatic heart failure, and LVEF $\leq 35\%$ despite >3 months of OMT	IIa	A
An ICD should be considered in patients with DCM with a genotype associated with high SCD risk and LVEF $>35\%$ in the presence of additional risk factors*	IIa	C
An ICD may be considered in selected patients with DCM with a genotype associated with high SCD risk and LVEF $>35\%$ without additional risk factors	IIb	C
An ICD may be considered in patients with DCM without a genotype associated with high SCD risk and LVEF $>35\%$ in the presence of additional risk factors*	IIb	C
Subcutaneous defibrillator should be considered as an alternative to transvenous defibrillator in patients with an indication for an ICD when pacing therapy for bradycardia, cardiac resynchronization, or ATP is not needed	IIa	B
A wearable ICD may be considered for patients with HF who are at risk of sudden cardiac death for a limited period or as a bridge to an implanted device	IIb	B

ICD, implantable cardiac defibrillator; DCM, dilated cardiomyopathy; OMT, optimal medical therapy; LVEF, left ventricular ejection fraction; SCD, sudden cardiac death; ATP, anti tachy pacing; HF, heart failure. *Additional risk factors include syncope, late gadolinium enhancement presence on cardiac magnetic resonance.

symptoms. However, the natural history of AC is different from that of DCM. Recovery in abstainers is associated with an excellent prognosis. For this reason, we recommended complete abstinence from alcohol to our patient, and we have seen an improvement in the EF over the follow-up period [12]. However, the identification of an acquired cause does not rule out an underlying genetic variant, while the latter, if present, may be triggered earlier when an acquired cause intervenes [2,8]. Drugs that are used for ADHD treatment may have potential adverse cardiovascular effects. These agents exert stimulant effects on the central nervous system, increase norepinephrine and dopamine levels in the prefrontal cortex and stimulate adrenergic receptors in the heart and blood. Sympathetic nervous system stimulation can lead to an increased risk of cardiovascular mortality [13]. Some of the drugs taken by the patient could be a further cause of heart failure, such as gabapentin. Other drugs may have exposed the patient to an increased risk of life-threatening arrhythmias or other cardiovascular events (myocardial infarction, Takotsubo syndrome, SCD) as in the case of quetiapine, trazodone and atomoxetine [3,13]. It has been reported that adults with DCM have an annual incidence of SCD between 2 and 4 percent [1,3]. According to recent ESC guidelines on cardiomyopathies, ICD implantation in primary prevention is indicated in patients symptomatic for heart failure and $FE \leq 35\%$, despite 3 months of optimized medical therapy, and in patients with $FE \geq 35\%$ and additional risk factors, for example, high arrhythmic risk genotype, arrhythmias, and late gadolinium enhancement on cardiac MRI (documented in our patient, Figure 1) (Table 1). The choice of ICD type depends on the clinical setting [7,14]. In the case of our patient, who underwent a diagnostic definition and was at risk of SCD, a WCD (class II-B) was suggested [14,15]. WCD allows managing the transient risk of arrhythmic death pending SCD risk reduction or permanent protection or definitive indication for ICD implantation [16]. Although WCD use does not have a class I recommend in the most recent guidelines, its use in clinical practice is growing [15-17]. Indeed, even if the guidelines provide a valuable reference for treatment choices, it is necessary to consider that the decision at the individual level may be influenced by several factors such as age, psychosocial conditions, and the patient's will, factors that are not always taken sufficiently into account by the guidelines. We planned to improve medication for 3 months and then re-examine the patient protected by the WCD.

In our case, the psychiatric condition made our management decision challenging. According to the psychiatrist who followed the patient during his hospital stay, adherence to treatment was considered a serious problem, both about taking medication at home and, in particular, about WCD. For this reason, the choice, shared with other specialists, fell on the definitive ICD [1,14,16]. The indication for S-ICD implantation was established based on the patient's age and possible long-term complications secondary to the implantation of intravascular leads in the case of an ICD (Table 1) [1,17]. The S-ICD has been proven to be a valuable alternative to transvenous defibrillators in all patients who do not require bradycardia pacing, resynchronization, or anti-tachycardia pacing (class II-B) [17,18]. As a matter of fact, an S-ICD is considered the best treatment option before the age of 50 [18,19]. We chose the S-ICD also because our patient could be at risk of endocarditis (history of drug addiction), and we did not know if he could undergo a transplantation or ventricular assist device (VAD) implantation that requires the protection of the vascular access site to limit the risk of systemic infection. Furthermore, the potential for explantation and the possibility of complications associated with the procedure must be considered. Therefore, in our patient, the use of a WCD would have allowed us to have a cardiac MRI not hindered by artifacts that limited examination interpretation (Figure 2) [15,20].

Conclusions

This case report emphasizes that in clinical practice, though the guidelines provide a valuable reference for treatment choices, it is necessary to consider that the decision at the individual level may be influenced by several factors, including age, psychosocial conditions, and the patient's will, that are not always taken into sufficient account by guidelines.

References

- Arbelo E, Protonotarios A, Gimeno JR, et al. 2023 ESC Guidelines for the management of cardiomyopathies. *Eur Heart J* 2023;44:3503-626.



2. Lei Z, Cao J, Wu J, Lu Y, et al. Identification of the communal pathogenesis and immune landscape between viral myocarditis and dilated cardiomyopathy. *ESC Heart Fail* 2024;11:282-92.
3. Orphanou N, Papatheodorou E, Anastakis A. Dilated cardiomyopathy in the era of precision medicine: latest concepts and developments. *Heart Fail Rev* 2022;27:1173-91.
4. Merlo M, Cannatà A, Gobbo M, et al. Evolving concepts in dilated cardiomyopathy. *Eur J Heart Fail* 2018;20:228-39.
5. William Bracamonte-Baran, Daniela Čiháková. Cardiac autoimmunity: myocarditis. *Adv Exp Med* 2017;1003:187-221.
6. Ekström K, Lehtonen J, Kandolin R, et al. Long-term outcome and its predictors in giant cell myocarditis. *Eur J Heart Fail* 2016;18:1452-8.
7. D'Ambrosio A, Patti G, Manzoli A, et al. The fate of acute myocarditis between spontaneous improvement and evolution to dilated cardiomyopathy: a review. *Heart* 2001;85:499-504.
8. Yamada T, Nomura S. Recent findings related to cardiomyopathy and genetics. *Int J Mol Sci* 2021;22:12522.
9. Cooper LT, Baughman KL, Feldman AM, et al. The role of endomyocardial biopsy in the management of cardiovascular disease: a scientific statement from the American Heart Association, the American College of Cardiology, and the European Society of Cardiology Endorsed by the Heart Failure Society of America and the Heart Failure Association of the European Society of Cardiology. *Eur Heart J* 2007;28:3076-93.
10. Könnemann H, Dagres N, Merino JL, et al. Spotlight on the 2022 ESC guideline management of ventricular arrhythmias and prevention of sudden cardiac death: 10 novel key aspects. *Europace* 2023;25:euad091.
11. Fernández-Solà J. The effects of ethanol on the heart: alcoholic cardiomyopathy. *Nutrients* 2020;22:572.
12. Domínguez F, Adler E, García-Pavía P. Alcoholic cardiomyopathy: an update. *Eur Heart J* 2024;45:2294-305.
13. Torres-Acosta N, O'Keefe JH, O'Keefe CL, Lavie CJ. Cardiovascular effects of ADHD therapies: JACC review topic of the week. *J Am Coll Cardiol* 2020;76:858-66.
14. Zeppenfeld K, Tfelt-Hansen J, de Riva M, et al. 2022 ESC Guidelines for the management of patients with ventricular arrhythmias and the prevention of sudden cardiac death. *Eur Heart J* 2022;43:3997-4126.
15. Casolo G, Gulizia MM, Aschieri D, et al. Position paper ANMCO: guida all'uso appropriato del defibrillatore indossabile nella pratica clinica per i pazienti ad elevato rischio transitorio di morte improvvisa. *G Ital Cardiol* 2023;24:394-411. [Article in Italian].
16. Mc Donagh TA, Metra M, Adamo M, et al. 2021 ESC guidelines for the diagnosis and treatment of acute and chronic heart failure. *Eur Heart J* 2021;42:3599-726.
17. Boriani G, Vitolo M, Leyva F. Implantable cardioverter-defibrillators for primary prevention of sudden cardiac death: what are the barriers to implementation in the 'real world'? *Eur J Heart Fail* 2022;24:1223-6.
18. Probst V, Boveda S, Sadoul N et al. Subcutaneous implantable cardioverter defibrillator indication in prevention of sudden cardiac death in difficult clinical situations: a French expert position paper. *Arch Cardiovasc Dis* 2020;113:359-66.
19. Migliore F, Mattesi G, Zorzi A, et al. Il defibrillatore sottocutaneo nella pratica clinica *G Ital Cardiol* 2019;20:641-50. [Article in Italian].
20. Holtstiege V, Meier C, Bietenbeck M, et al. Clinical experience regarding safety and diagnostic value of cardiovascular magnetic resonance in patients with a subcutaneous implanted cardioverter/defibrillator (S-ICD) at 1.5 T. *J Cardiovasc Magn Reson* 2020;22:35.

Received: 6 August 2024; Accepted: 30 January 2025; Early view: 1 April 2025.

Contributions: all the authors made a substantive intellectual contribution, have read and approved the final version of the manuscript.

Conflict of interest: the authors declare that they have no competing interests.

Ethics approval and consent to participate: institutional review board approval was not required for this study as only de-identified compliant data were used in the analysis. The Data Protection Act and the Helsinki Declaration's principles were followed in the study.

Patient consent for publication: the patient gave his written consent to use his personal data for the publication of this case report and any accompanying images.

Availability of data and materials: all mentioned data in the paper are available in the medical records.

Publisher's note: all claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).

