

©Copyright: the Author(s), 2024  
Licensee PAGEPress, Italy

# Evaluation of asthma control after reinforcement of proper inhaler techniques in a tertiary care center in northern India

Manav Raj Singh Gill, Sparsh Gupta, Amit Sharma, Pranav Ish, Deepak Pandey

Department of Pharmacology, Vardhman Mahavir Medical College, Safdarjung Hospital, New Delhi, India

## Abstract

Asthma is an obstructive lung disease with high prevalence. Symptom control in asthma remains poor despite significant advancements in treatment guidelines and inhalational devices. This is often attributed to inadequate patient adherence to the inhaler technique and insufficient education on its long-term use. Through this study, we aimed to assess the impact of reinforcement of proper inhaler technique on asthma symptom control and quality of life. A prospective cohort of 400 asthma patients using dry powder inhalers and metered dose inhalers was recruited in the outpatient clinic of our hospital. At the time of recruitment, the correctness of inhaler technique usage was assessed, and the pre-test Inhaler Technique Score (ITS) was recorded for each patient. Then, a ten-step inhalation technique was taught through demonstration, and the post-test ITS score was recorded again. This was done for each patient at monthly intervals for 3 months. Along with it, two tailored questionnaires, the Asthma Control Test (ACT) and Asthma Quality of Life Questionnaire (AQLQ), were administered at each visit to assess disease control and quality of life. Significant enhancements in inhalation technique were noted from the first to the third visit using the ten-step inhalation usage scores, which improved from 6.91 to 9.87 (out of 10) ( $p < 0.001$ ), paralleled by ACT score increases from 17.06 to 19.37 (out of 25) ( $p < 0.001$ ) from visits one to two, and from 19.37 to 20.52 (out of 25) ( $p < 0.001$ ) from visits two to three, signifying improvement in symptom control. Quality of life similarly improved from 4.45 to 5.12 to 5.45 (out of 7) ( $p < 0.001$ ) across the three visits as assessed by AQLQ. In conclusion, promoting proper inhaler technique through structured education programs is crucial for optimizing long-term asthma management and enhancing patients' quality of life.

**Key words:** inhaler technique, metered dose inhaler, dry powder inhaler, adherence, quality-of-life, patient education.

Correspondence to: Sparsh Gupta, Department of Pharmacology, VMMC and Safdarjung Hospital, New Delhi -110029, India.  
Tel.: +91-0-9818057272. E-mail: healing\_sparsh@yahoo.co.in

## Introduction

Inhalational medications are the mainstay in contemporary asthma treatment due to their ability to deliver therapeutic doses directly to the lungs while minimizing systemic side effects [1]. Effective inhaler use is critical for achieving optimal symptom control and disease management. However, ineffective techniques can compromise drug delivery, leading to decreased efficacy and poor outcomes in asthma management. Patient adherence to prescribed inhalation therapies significantly influences treatment effectiveness [2-4]. Studies have highlighted that a substantial proportion of patients, over 60% in some reports, do not use their inhalers correctly, thereby reducing the medication's efficacy [5,6]. This underscores the importance of patients' knowledge of inhaler technique as well as the ability of healthcare providers to instruct on proper technique on the treatment outcomes [7,8].

While inhaler devices vary in dose preparation and operational mechanics, fundamental principles of correct inhalation technique apply universally. Hence, personalized patient education, assessment, and training are crucial components of effective

asthma care [9,10]. Guidelines universally recommend that patients should not start using a new inhaler device without adequate training from a healthcare professional and demonstration of correct technique [11].

Despite the clear guidelines advocating for proper inhaler technique training, there remains a scarcity of qualitative data assessing asthma control following reinforcement of correct inhaler technique. Thus, this study was carried out to assess patient compliance with the correct inhaler technique and the effect of training interventions on the same, thereby serving as a guide to improved clinical practice and patient management strategies.

## Materials and Methods

### Study population

The study was conducted in the Department of Pharmacology and the Department of Pulmonology in a tertiary care center in Northern India. It included 400 asthma patients observed over monthly visits for 3 months. The sample size was determined based on Singh *et al.* [12]. Inclusion criteria were



age >18 years and use of inhaler therapy for at least 3 months. Exclusion criteria included participation in other clinical trials, terminal illness, and initiation of new asthma therapy during the study.

## Study design

The study was a prospective cohort study conducted for a duration of 12 months from August 2022 to August 2023. The objective was to assess the compliance of patients to proper inhaler technique through an inhaler technique score (ITS), which ranged from 0 to 10 (*Supplementary Table 1*), with a score of 0 indicating no compliance to 10 indicating full compliance. Then, the reinforcement of the proper inhaler technique was done *via* a 10-step technique demonstration, and its effect on the ITS score was evaluated. Further evaluation was done to analyze the effect of the correct inhaler technique on asthma control using the Asthma Control Test (ACT). The ACT scores range from 5 (poor control) to 25 (complete control), with scores >19 indicating well-controlled asthma. The change in quality of life post-reinforcement with the Asthma Quality of Life Questionnaire (AQLQ) was also observed. AQLQ is rated on a 7-point Likert scale, where 1 denotes maximal impairment and 7 denotes no impairment in quality of life. The study procedure is summarized as follows.

## Baseline

Participants completed a pretest on inhaler technique knowledge (ITS), ACT, and AQLQ. Following a demonstration of inhaler technique steps, a post-test ITS score was obtained.

## Week 4

Participants were reassessed using ACT and AQLQ. A second ITS demonstration was given as reinforcement, with ITS scores recorded before the demonstration.

## Week 8

Final assessment was conducted using ITS, ACT, and AQLQ.

## Results

Out of the 400 patients, there was no dropout, and the data of all patients were recorded and analyzed. The average values of demographic parameters and anthropometric details of the study participants (n=400) are shown in Table 1.

The mean ITS pre-test scores (out of 10) of all patients at baseline, week 4, and week 8 were  $6.9\pm 1.7$ ,  $9.55\pm 1.017$ , and  $9.87\pm 0.719$ , respectively. A paired *t*-test was done to compare ITS score values between baseline and week 4, and further between week 4 and week 8. There was a significant increase in the ITS score from baseline to week 4 ( $p<0.05$ ). Similarly, there was a significant increase in the ITS score from week 4 to week 8 ( $p<0.05$ ). The mean post-test ITS score at each monthly visit for all patients came out to be 10.

The ACT scores were also recorded for each patient at baseline, week 4, and week 8. The mean ACT score (out of 25) for all patients came out to be  $17.1\pm 3.3$ ,  $19.4\pm 2.7$ , and  $20.5\pm 2.3$  at baseline, week 4, and week 8, respectively. On paired *t*-tests, there was a significant ( $p<0.05$ ) increase in ACT values from baseline to week 4 and further from week 4 to week 8.

The AQLQ scores were also recorded for each patient at baseline, week 4, and week 8. The mean AQLQ score (out of 7) for all patients came out to be  $4.45\pm 1.03$ ,  $5.12\pm 0.85$ , and  $5.45\pm 0.73$  at baseline, week 4, and week 8, respectively. On paired *t*-tests, there was a significant ( $p<0.05$ ) increase in AQLQ values from baseline to week 4 and further from week 4 to week 8.

A Pearson correlation analysis was performed to evaluate the role of the ITS score in influencing the ACT score and AQLQ score. There was a significant positive correlation ( $r^2=0.659$ ;  $p<0.05$ ) between the change in ITS scores from baseline to week 4 and the change in ACT scores from baseline to week 4. Similarly, there was a significant positive correlation ( $r^2=0.539$ ;  $p<0.05$ ) between the change in ITS scores from week 4 to week 8 and the change in ACT scores from week 4 to week 8. Likewise, there was a significant positive correlation ( $r^2=0.583$ ;

**Table 1.** Demographic and clinical parameters of all patients at baseline, 4 weeks, and 8 weeks.

	Baseline	4 weeks	8 weeks
Age (median)	38 years		
Gender (male)	43.8%		
Weight (kg)	63.4±13.3		
Height (m)	1.6±0.08		
BMI (kg/m <sup>2</sup> )	24.4±4.7		
Pretest ITS score <sup>a</sup> (mean±SD)	6.91±1.71	9.55±1.01*	9.87±0.71**
Post-test ITS score <sup>a</sup> (mean±SD)	10	10	10
ACT score <sup>b</sup> (mean±SD)	17.06±3.27	19.37±2.67 *	20.52±2.31**
AQLQ score <sup>c</sup> (mean±SD)	4.45±1.03	5.12±0.85 *	5.45±0.73**

BMI, body mass index; ITS, inhaler technique score; SD, standard deviation; ACT, asthma control test; AQLQ, asthma quality of life questionnaire; <sup>a</sup>total score out of 10; <sup>b</sup>total score out of 25; <sup>c</sup>total score out of 7; \* $p<0.05$  on paired *t*-test between scores of weeks 4 and day 0; \*\* $p<0.05$  on paired *t*-test between scores of week 8 and week 4.



$p < 0.05$ ) between the change in ITS scores from baseline to week 4 and the change in AQLQ scores from baseline to week 4. Also, there was a significant positive correlation ( $r^2 = 0.404$ ;  $p < 0.05$ ) between the change in ITS scores from week 4 to week 8 and the change in AQLQ scores from week 4 to week 8.

## Discussion

The pre-test ITS score at baseline was conducted amongst the participants who were already on inhalers for a minimum duration of the past 3 months. The results had a mean score of  $6.91 \pm 1.71$ , with a maximum score of 10 and a minimum score of 4. This emphasizes the very basis of this study that, despite timely diagnosis and proper medication, patients are not following the correct inhaler technique. Our findings are consistent with an earlier study by Lia Jehedi *et al.* that found that even though most asthmatic patients were confident in their technique, many did not use their inhaler(s) correctly [13]. The pre-test ITS score at week 4 had a mean value of  $9.55 \pm 1.01$ , with significant improvement ( $p < 0.05$ ) from baseline values. This indicates that if the participants are taught the inhaler technique along with a proper demonstration of each step in the correct sequence, their compliance with the inhaler technique improves. Further, the pre-test ITS score at week 8 had a mean value of  $9.87 \pm 0.71$  with significant improvement over the ITS score values at week 4. This shows that the reinforcement of the inhaler technique can be an essential intervention for the continued success of the treatment over time. These results demonstrate the crucial role of proper inhaler technique in the management of asthma [13]. The post-test ITS scores demonstrating short-term learning of the patient on all three occasions were found to be 10 in all 400 participants. This was because it was ensured that every patient should learn the inhaler technique correctly before leaving the outpatient department, and to ensure that patients would follow the steps accordingly while taking the medication.

The ACT was done to assess the level of symptom control of asthma in the patient at baseline, week 4, and week 8. The mean value of ACT at baseline was  $17.06 \pm 3.27$ , which means that asthma was not well controlled. This significantly ( $p < 0.05$ ) improved ACT scores, being  $19.37 \pm 2.67$  and  $20.52 \pm 2.31$  at week 4 and week 8, respectively, indicating a transition to well-controlled asthma. On Pearson's correlation, we found out that the change in pre-test ITS score from one visit to the subsequent visit significantly correlated with the corresponding change in ACT scores. This highlights the impact of adherence to the correct inhaler technique on symptom control in asthma. In previous studies, it has been similarly found that the adherence of patients to medications improved disease control and decreased the risk of exacerbations as well [14-16].

Asthma can be a crippling disorder that negatively impacts a person's quality of life. In this study, the AQLQ score was calculated for each patient at baseline, week 4, and week 8 to assess their quality of life. The mean value of AQLQ at baseline was  $4.45 \pm 1.03$ , which means that there was moderate limitation, which significantly improved to  $5.12 \pm 0.85$  and  $5.45 \pm 0.73$  at week 4 and week 8, respectively. On Pearson's correlation, the change in AQLQ values from baseline to week 4 correlated significantly with the corresponding change in pre-test ITS scores and ACT scores from baseline to week 4. Similarly, the change in AQLQ values from week 4 to week 8 also correlated significantly with the

corresponding change in pre-test ITS scores and ACT scores from week 4 to week 8 [17]. This becomes significant for asthma patients because they must keep taking medication for longer durations, and the quality of life is an apt measure to evaluate the overall treatment effectiveness.

## Conclusions

The reinforcement of proper inhaler technique results in better asthma control and better quality of life in patients with bronchial asthma. This places the healthcare provider in a crucial position to oversee inhaler technique and interact with the patient to enhance adherence. Also, simple tools like demonstration of inhaler technique steps and checklists can have a major impact on adherence, thus leading to better treatment outcomes.

## References

1. Boulet LP, Reddel HK, Bateman E, et al. The global initiative for asthma (GINA): 25 years later. *Eur Respir J* 2019;54:1900598.
2. Virchow JC, Crompton GK, Dal Negro R, et al. Importance of inhaler devices in the management of airway disease. *Resp Med* 2008;102:10-9.
3. Bousquet J, Mantzouranis E, Cruz AA, et al. Uniform definition of asthma severity, control, and exacerbations: document presented for the World Health Organization consultation on severe asthma. *J Allergy Clin Immunol* 2010;126:926-38.
4. Westerik JA, Carter V, Chrystyn H, et al. Characteristics of patients making serious inhaler errors with a dry powder inhaler and association with asthma-related events in a primary care setting. *J Asthma* 2016;53:321-9.
5. Melani AS, Bonavia M, Cilenti V, et al. Inhaler mishandling remains common in real life and is associated with reduced disease control. *Resp Med* 2011;105:930-8.
6. Press VG, Arora VM, Shah LM, et al. Misuse of respiratory inhalers in hospitalized patients with asthma or COPD. *J Gen Intern Med* 2011;26:635-42.
7. Van der Palen J, Klein JJ, Kerkhoff AH, et al. Evaluation of the effectiveness of four different inhalers in patients with chronic obstructive pulmonary disease. *Thorax* 1995;50:1183-7.
8. Lavorini F, Magnan A, Dubus JC, et al. Effect of incorrect use of dry powder inhalers on management of patients with asthma and COPD. *Resp Med* 2008;102:593-604.
9. Dekhuijzen PN, Vincken W, Virchow JC, et al. Prescription of inhalers in asthma and COPD: towards a rational, rapid and effective approach. *Resp Med* 2013;107:1817-21.
10. Price D, Bosnic-Anticevich S, Briggs A, et al. Inhaler competence in asthma: common errors, barriers to use and recommended solutions. *Resp Med* 2013;107:37-46.
11. Vestbo J, Hurd SS, Agustí AG, et al. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. *Am J Resp Crit Care Med* 2013;187:347-65.
12. Singh S, Salvi S, Mangal DK, et al. Prevalence, time trends and treatment practices of asthma in India: the Global Asthma Network study. *ERJ Open Res* 2022;8:00528-2021.
13. Jahedi L, Downie SR, Saini B, et al. Inhaler technique in asthma: how does it relate to patients' preferences and attitudes



- toward their inhalers? *J Aerosol Med Pulm Drug Deliv* 2017; 30:42-52.
14. Beuther DA, Sutherland ER. Overweight, obesity, and incident asthma: a meta-analysis of prospective epidemiologic studies. *Am J Resp Crit Care Med* 2007;175:661-6.
  15. Scullion J. The nurse practitioners' perspective on inhaler education in asthma and chronic obstructive pulmonary disease. *Can Respir J* 2018;2018:2525319.
  16. Schatz M, Sorkness CA, Li J, et al. Asthma control test: reliability, validity, and responsiveness in patients not previously followed by asthma specialists. *J Allergy Clin Immunol* 2006; 117:549-56.
  17. Juniper EF, Buist AS, Cox FM, et al. Validation of a standardized version of the asthma quality of life questionnaire. *Chest* 1999;115:1265-70.

---

*Online supplementary material:*

*Supplementary Table 1. Inhalation technique steps and score.*

---

Received: 11 September 2024; Accepted: 13 September 2024; Early view: 21 October 2024.

Contributions: all authors were involved in the literature search, planning, conduct, writing the original draft of the manuscript, literature search, and editing of the study. All the authors have agreed with the submitted manuscript.

Conflict of interest: the authors declare that they have no conflicts of interest regarding the current study.

Ethics approval and consent to participate: ethical clearance obtained from the Institutional Ethics Committee, VMMC, and Safdarjung Hospital, New Delhi, with serial no IEC/VMMC/SJH/Thesis/06/2022/CC-240 dated 11th July 2022.

Informed consent: written informed consent was taken from all participants.

Patient consent for publication: written informed consent has been obtained from all patients for the publication of the data.

Availability of data and materials: data and materials are available from the corresponding author upon request.

*Publisher's note: all claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*

*This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).*

