doi: 10.4081/monaldi.2025.3354

SUPPLEMENTARY MATERIAL

Histopathological assessment of thymoma based on the World Health Organization 2021 classification with emphasis on transcapsular invasion: experience from a tertiary care center

Saloni,[1] Renuka Malipatel,[2] Sreekar Balasundaram[3]

[1]Department of Pathology, Guru Gobind Singh Medical College, Faridkot, Punjab; [2]Department of Pathology, St John's Medical College and Hospital, Bangalore, Karnataka; [3]Department of Cardiothoracic Surgery, St John's Medical College and Hospital, Bangalore, Karnataka, India

Correspondence: Saloni, Department of Pathology, Guru Gobind Singh Medical College, Vijay hospital near durga mandir Giddarbaha, Sri Muktsar Sahib, Faridkot, Punjab, 152101, India. Tel.: +9194633341830. E-mail: saloni.bansal4@gmail.com

Key words: histological subtype, morphology, prognosis, staging, capsular invasion.



Supplementary Figure 1. Histological subtypes, morphological criteria as per the World Health Organization 2021 classification of thymoma.



Supplementary Figure 2. A) Association of clinical stage with histological subtype; B) outcome distribution; C) survival analysis for stage I and stage II patients.



Supplementary Figure 3. Mixed subtypes of thymoma with varying proportions of B2 and B3 subtypes. CK 19 highlights denser meshwork of thymic epithelial cells in B3 areas (right side of the field, depicted by the arrowhead) as compared to B2 areas (left side of the field, pointed by arrow); Tdt shows lesser density of lymphocytes in B3 areas (100× magnification).