

## Authors' Response

Branko Beronja,<sup>1</sup> Aleksandra Karan,<sup>2</sup> Biljana Lukic,<sup>2</sup> Ivana Milosevic,<sup>1,3</sup> Jelena Dotlic,<sup>1,4</sup> Tatjana Gazibara<sup>5</sup>

<sup>1</sup>Faculty of Medicine, University of Belgrade; <sup>2</sup>General Hospital "Dr Radivoj Simonovic", Sombor; <sup>3</sup>Clinic of Infectious and Tropical Diseases, University Clinical Center of Serbia, Belgrade; <sup>4</sup>Clinic of Obstetrics and Gynecology, University Clinical Center of Serbia, Belgrade; <sup>5</sup>Institute of Epidemiology, Faculty of Medicine, University of Belgrade, Serbia

**Key words:** COVID-19, smoking, mortality.

Correspondence to: Tatjana Gazibara, Institute of Epidemiology, Faculty of Medicine University of Belgrade, Visegradska 26a, 11000 Belgrade, Serbia. Tel.: +381-11-3607-062. E-mail: tatjana.gazibara@med.bg.ac.rs

Dear Editor,

We read the comments by Rossato and Di Vincenzo with enthusiasm and appreciate their interest in our study, which helps deepen the discussion on smoking and COVID-19 [1]. Indeed, smoking is a major health issue worldwide, as it is associated with a variety of poor outcomes, including COVID-19. Because of the widespread tobacco use in Serbia (36.6%) [2], it is estimated that male smokers live 15 years shorter, and women smokers live 17 years shorter than their non-smoking age-matched counterparts [3]. Furthermore, research on household budget suggests that smokers in Serbia would rather spend money on cigarettes than on food, clothes and education [4].

That being said, the prevalence of smokers among hospitalized patients with COVID-19 in the Beronja *et al.* study (40.7% ever smokers vs. 59.3% non-smokers) is even slightly higher than the prevalence of smoking in the general population in Serbia [2,5]. Nevertheless, it would be unrealistic to expect that smokers make up the majority of hospitalized patients with COVID-19, because there are almost twice as many non-smokers in the Serbian general population and smokers may be less susceptible to catch COVID-19 [6]. While data from one center, such as that from the Beronja *et al.* study [5], cannot be generalized to the entire country, they are suggestive of the distribution of smokers and non-smokers in the Serbian population. This high prevalence of tobacco consumption requires more intense and systematic structural measures of smoking prevention and cessation, but these aspects fall outside of the scope of the Beronja *et al.* study [5].

In the study by Beronja *et al.* [5], women accounted for 68.7% of patients treated for severe COVID-19 in the hospital setting. This result was certainly unexpected, given that the body of literature suggests that men are at risk of poorer COVID-19 outcomes [7]. It should be kept in mind that women in Serbia make up the majority of informal (*i.e.*, unpaid) caregivers of family members, particularly in multi-generational households, and were therefore, more exposed to COVID-19 [8]. This burden associated with caregiving can lead to general health deterioration, which may be associated with greater needs for hospital attention when infected with COVID-19 [9]. An analysis of more than one million people in Canada suggests that women are especially susceptible to COVID-19 during peaks of epidemic waves over the first two pandemic years and at those times

the proportion of women with COVID-19 exceeded that of men [10]. This is, no doubt, intriguing and merits further exploration especially through the lens of gender inequity.

Pre-existing chronic illnesses of hospitalized COVID-19 patients were proven critical for understanding of COVID-19 prognosis and outcomes in different studies [1,11], including a study conducted in the Serbian population, where diabetes, neurological disorders, and recent injuries/fractures were associated with a higher risk of respiratory failure due to COVID-19 [12]. For this reason, in the study of Beronja *et al.* [5], major chronic illnesses were accounted for. Out of those, the presence of respiratory illnesses and obesity were associated with poorer COVID-19 outcomes among smokers. However, none of the pre-existing chronic illnesses were predictive of poorer survival in non-smokers hospitalized with COVID-19 [5]. Thus, studies focusing on interactions between nicotine and components of the immune system among people with and without chronic illnesses who smoke are necessary to identify the underlying mechanisms of how tobacco prevents the onset of the infection, but conversely, contributes to deterioration once a smoker becomes infected.

Effects of smoking on different aspects of health must not be underestimated. Smoking behaviors should be continuously scrutinized and recorded. Continuous education campaigns about detrimental effects of tobacco are needed not only in prevention of chronic non-communicable diseases, but also in relation to outcomes of acute infectious diseases.

## References

1. Rossato M, Di Vincenzo A. Comments on "Smoking patterns and outcomes of severe SARS-CoV-2 infection: a retrospective cohort study". Monaldi Arch Chest Dis 2025; doi: 10.4081/monaldi.2025.3430.
2. Global Action to End Smoking. State of Smoking and Health in Serbia. 2024. Available from: <https://globalactiontoendsmoking.org/research/tobacco-around-the-world/serbia/>.
3. Kilibarda B, Vasic M, Gudelj Rakic, Atanasijevic D. Smoking-attributable life and working years lost in Serbia. Tob Prev Cessation 2023;9:A167.
4. Vladisavljevic M, Zubovic J, Jovanovic O, Đukić M.



- Crowding-out effect of tobacco consumption in Serbia. *Tob Control* 2024;33:s88-94.
5. Beronja B, Karan A, Lukic B, et al. Smoking patterns and outcomes of severe SARS-CoV-2 infection: a retrospective cohort study. *Monaldi Arch Chest Dis* 2025; doi: 10.4081/monaldi.2025.2916.
  6. Paleiron N, Mayet A, Marbac V, et al. Impact of tobacco smoking on the risk of COVID-19: a large scale retrospective cohort study. *Nicotine Tob Res* 2021;23:1398-404.
  7. Patanavanich R, Siripoon T, Amponnavarat S, Glantz SA. Active smokers are at higher risk of COVID-19 death: a systematic review and meta-analysis. *Nicotine Tob Res* 2023;25:177-84.
  8. Todorovic N, Vracevic M, Rajovic N, et al. Quality of life of informal caregivers behind the scene of the COVID-19 epidemic in Serbia. *Medicina* 2020;56:647.
  9. Del Río-Lozano M, García-Calvente M, Elizalde-Sagardia B, Maroto-Navarro G. Caregiving and caregiver health 1 year into the COVID-19 pandemic (CUIDAR-SE Study): a gender analysis. *Int J Environ Res Public Health* 2022;19:1653.
  10. Wu C, Qian Y. The gender peak effect: women are most vulnerable to infections during COVID-19 peaks. *Front Public Health* 2022;10:937179.
  11. Emami A, Javanmardi F, Pirbonyeh N, et al. Prevalence of underlying diseases in hospitalized patients with COVID-19: a systematic review and meta-analysis. *Arch Acad Emerg Med* 2020;8:e35.
  12. Beronja B, Gazibara T, Dotlic J, et al. Pre-existing chronic illnesses as a risk factor for the onset of respiratory failure due to COVID-19. *Clin Epidemiol Glob Health* 2024;29:101756.

---

Received: 5 March 2025; Accepted: 14 March 2025; Early view: 17 March 2025.

Contributions: Branko Beronja, Jelena Dotlic, Tatjana Gazibara: draft of the manuscript. Aleksandra Karan, Biljana Lukic, Ivana Milosevic: critical review of the manuscript.

Conflict of interest: the authors declare that they do not have potential conflicts of interest.

Ethics approval and consent to participate: not applicable.

Informed consent: not applicable.

Patient consent for publication: not applicable.

Availability of data and materials: not applicable.

*Publisher's note: all claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*

*This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).*

