

SUPPLEMENTARY MATERIAL


To evaluate the quit rate in smoking and smokeless tobacco users accessing the National Tobacco Quitline Services, New Delhi

Raj Kumar,^{1,2} Ritika Gupta,² Praveen Sinha,³ Shyam Dubey,² Anil Kumar Mavi⁴


¹Department of Pulmonary Medicine, Vallabhbhai Patel Chest Institute, University of Delhi; ²National Tobacco Quit Line Services, Vallabhbhai Patel Chest Institute, University of Delhi; ³Tobacco Control, World Health Organization Country Office, New Delhi; ⁴Department of Botany and Life Sciences, Sri Aurobindo College, University of Delhi, Delhi, India

Correspondence: Raj Kumar, Department of Pulmonary Medicine, Vallabhbhai Patel Chest Institute, University of Delhi, Delhi-110007, India. E-mail: rajkumarvpai@gmail.com


Key words: tobacco quit line, smoking tobacco, smokeless tobacco, environment tobacco smoke.




Supplementary Figure 1. Overview of quit rates in registered tobacco users.




Supplementary Figure 2. Quitter and registered category wise.




Supplementary Figure 3. Smokeless tobacco age distribution.




Supplementary Figure 4. Smoking tobacco age distribution.




Supplementary Figure 5. Smokeless tobacco education distribution.




Supplementary Figure 6. Smoking education distribution.



Supplementary Figure 7. Income distribution.



Supplementary Figure 8. Tobacco usage by number of years of use.



Supplementary Figure 9. Tobacco users by quantity of tobacco used per day.